1895

Shorter Contributions to Paleontology and Stratigraphy

U.S. GEOLOGICAL SURVEY BULLETIN 1895

I 19.3:

best copy available

AVAILABILITY OF BOOKS AND MAPS OF THE U.S. GEOLOGICAL SURVEY

Instructions on ordering publications of the U.S. Geological Survey, along with prices of the last offerings, are given in the current-year issues of the monthly catalog "New Publications of the U.S. Geological Survey." Prices of available U.S. Geological Survey publications released prior to the current year are listed in the most recent annual "Price and Availability List." Publications that are listed in various U.S. Geological Survey catalogs (**see back inside cover**) but not listed in the most recent annual "Price and Availability List" are no longer available.

Prices of reports released to the open files are given in the listing "U.S. Geological Survey Open-File Reports," updated monthly, which is for sale in microfiche by the U.S. Geological Survey, Books and Open-File Reports Section, Federal Center, Box 25425, Denver, CO 80225. Reports released through the NTIS may be obtained by writing to the National Technical Information Service, U.S. Department of Commerce, Springfield, VA 22161; please include NTIS report number with inquiry.

Order U.S. Geological Survey publications **by mail** or **over the counter** from the offices given below.

BY MAIL

Books

Professional Papers, Bulletins, Water-Supply Papers, Techniques of Water-Resources Investigations, Circulars, publications of general interest (such as leaflets, pamphlets, booklets), single copies of Earthquakes & Volcanoes, Preliminary Determination of Epicenters, and some miscellaneous reports, including some of the foregoing series that have gone out of print at the Superintendent of Documents, are obtainable by mail from

U.S. Geological Survey, Books and Open-File Reports
Federal Center, Box 25425
Denver, CO 80225

Subscriptions to periodicals (Earthquakes & Volcanoes and Preliminary Determination of Epicenters) can be obtained ONLY from the

Superintendent of Documents
Government Printing Office
Washington, D.C. 20402

(Check or money order must be payable to Superintendent of Documents.)

OVER THE COUNTER

Books

Books of the U.S. Geological Survey are available over the counter at the following U.S. Geological Survey Public Inquiries Offices, all of which are authorized agents of the Superintendent of Documents:

- WASHINGTON, D.C.—Main Interior Bldg., 2600 corridor, 18th and C Sts., NW.
- DENVER, Colorado—Federal Bldg., Rm. 169, 1961 Stout St.
- LOS ANGELES, California—Federal Bldg., Rm. 7638, 300 N. Los Angeles St.
- MENLO PARK, California—Bldg. 3, Rm. 4522, 345 Middlefield Rd.
- RESTON, Virginia—503 National Center, Rm. 12201, 12201 Sunrise Valley Dr.
- SALT LAKE CITY, Utah—Federal Bldg., Rm. 8105, 125 South State St.
- SAN FRANCISCO, California—Customhouse, Rm. 504, 555 Battery St.
- SPOKANE, Washington—U.S. Courthouse, Rm. 678, West 920 Riverside Ave.
- ANCHORAGE, Alaska—Rm. 101, 4230 University Dr.
- ANCHORAGE, Alaska—Federal Bldg., Rm. E-146, 701 C St.

Maps

For maps, address mail orders to:

U.S. Geological Survey, Map Distribution
Federal Center, Box 25286
Denver, CO 80225

Residents of Alaska may order maps from:

Alaska Distribution Section, U.S. Geological Survey
New Federal Building - Box 12
101 Twelfth Ave., Fairbanks, AK 99701

Maps

Maps may be purchased over the counter at the U.S. Geological Survey offices where books are sold (all addresses in above list) and at the following U.S. Geological Survey offices:

- ROLLA, Missouri—1400 Independence Rd.
- DENVER, Colorado—Map Distribution, Bldg. 810, Federal Center
- FAIRBANKS, Alaska—New Federal Bldg., 101 Twelfth Ave.

Shorter Contributions to Paleontology and Stratigraphy

Edited by WILLIAM J. SANDO

U.S. GEOLOGICAL SURVEY BULLETIN 1895

U.S. DEPARTMENT OF THE INTERIOR
MANUEL LUJAN, JR., Secretary

U.S. GEOLOGICAL SURVEY
Dallas L. Peck, Director

UNITED STATES GOVERNMENT PRINTING OFFICE: 1990

For sale by the
Books and Open-File Reports Section
U.S. Geological Survey
Federal Center, Box 25425, Denver, CO 80225

Library of Congress Cataloging in Publication Data

Shorter contributions to paleontology and stratigraphy
edited by William J. Sando
p. cm.—(U.S. Geological Survey bulletin ; 1895)
Includes bibliographical references
Supt. of Docs. no.: I 19.3: 1895
1. Paleontology. 2. Geology, stratigraphic. I. Sando,
William Jasper. II. Series.
QE75.B9 no. 1895
[QE721]
557.3 s—dc20 89-600217
[560] CIP

CONTENTS

CHAPTER A
Conodont Biostratigraphy of the Permian Road Canyon Formation, Glass Mountains, Texas
Bruce R. Wardlaw and Richard E. Grant

CHAPTER B
The Rugose Coral *Ankhelasma*—Index to Visean (Lower Carboniferous) Shelf Margin in the Western Interior of North America
William J. Sando, E. Wayne Bamber, and Barry C. Richards

CHAPTER C
Lithostratigraphy and Conodont Biostratigraphy Across the Lower-Middle Ordovician Disconformity (Early to Middle Whiterockian) at Pratt Ferry, Central Alabama
T.H. Shaw, Keith E. Roberson, and Anita G. Harris

Chapter A

Conodont Biostratigraphy of the Permian Road Canyon Formation, Glass Mountains, Texas

By BRUCE R. WARDLAW and RICHARD E. GRANT

U.S. GEOLOGICAL SURVEY BULLETIN 1895

SHORTER CONTRIBUTIONS TO PALEONTOLOGY AND STRATIGRAPHY

CONTENTS

Abstract A1
Introduction A1
The Type Section (RC I) A1
Supplementary Sections A3
The Leonardian-Guadalupian Boundary A4
Depositional Setting A4
Conodont Biofacies A5
Scope of Study A6
Genus *Hindeodus* A6
Genus *Neogondolella* A6
Genus *Neostreptognathodus* A7
Genus *Sweetina* A7
Genus *Sweetognathus* A8
Genus *Xaniognathus* A8
Measured Sections A11
References A16

PLATES

Plates follow References

1. *Neogondolella*
2. *Hindeodus* and *Neogondolella*
3. *Neostreptognathodus*, *Hindeodus*, *Sweetognathus*, and *Sweetina*
4. *Xaniognathus*, *Neostreptognathodus*, and *Neogondolella*

FIGURES

1. Map of southwestern portion of the Glass Mountains showing section locations of the Road Canyon Formation A2
2. Map of paleogeographic setting of the Permian of West Texas showing location of the Glass and northern Del Norte Mountains A4
3. Columnar sections of the Road Canyon Formation, Sections RC IX, RC IV, RC III, RC VI, RC V, and RC II A12
4. Columnar sections of the Road Canyon Formation, Sections RC I, RC VII, RC X, RC XI, RC VIII, and Rc. VII A13

TABLES

1. HOS-NOX ratio and percent of samples that yielded conodonts from sections of the Road Canyon Formation A5
2. Occurrence of conodont species in samples of the Road Canyon Formation sections that include underlying and overlying units A9

Conodont Biostratigraphy of the Permian Road Canyon Formation, Glass Mountains, Texas

By Bruce R. Wardlaw[1] and Richard E. Grant[2]

Abstract

The Road Canyon Formation is a shallow-water carbonate that was deposited on the seaward side of the narrow southern shelf of the Permian basin of West Texas. The conodont faunal succession of *Neogondolella idahoensis* to *Neogondolella serrata* generally occurs as a sharp transition in the lower part of the Road Canyon Formation. This faunal changeover is rapid and widespread. The Leonardian-Guadalupian Series boundary can be recognized as the level at which *N. idahoensis* is replaced by *N. serrata*.

INTRODUCTION

The unit now known as the Road Canyon Formation has undergone a slow but steady rise in stratigraphic status, beginning as a numbered member of a formation and ending up as the basis for a worldwide stage of the Permian. King (1931) established the unit as the "First Limestone member" of the Word Formation in the Glass Mountains of Texas, thus making it the lowermost stratigraphic unit of the Guadalupian Series. In keeping with a policy of naming the numbered units in the Glass Mountains, Cooper and Grant (1964) designated the "First Limestone member" as the Road Canyon Member of the Word Formation. Continued study of the brachiopods and the discovery of the ammonoid *Perrinites* in the Road Canyon led Cooper and Grant (1966) to the conclusion that the unit belonged more properly to the Leonardian Series and deserved formation status. However, the Road Canyon also contains the fusulinid *Parafusulina roth*, long considered to be a strong indicator of a Guadalupian age (Wilde, 1975). Because fusulinid workers did not agree that the Road Canyon was Leonardian, the age and correlations of the unit became controversial. As the controversy revolved about its assignment to either the Leonardian or Wordian Stage, one solution was to remove the unit from both and create a new Roadian Stage (Furnish, 1966, 1973). This solution retained the Leonardian as a stage, rather than as a series, and placed the Roadian in the Lower Permian (Furnish, 1973).

Wilde (1975) noted that, if only the Road Canyon and the correlative Cutoff Shale of the Apache Mountains were involved, Furnish's suggestion would be a simple and satisfactory solution to the apparent Road Canyon age anomaly. Unfortunately, other units are correlated to all or part of the Road Canyon, and complexities abound. Moreover, recognizing the Roadian as a stage implies potential worldwide usage, and, with the present bipartite subdivision of the Permian, the question remains: "Does the Roadian Stage belong in the Lower or the Upper Permian?" In addition to the problems bearing directly on the Road Canyon Formation, the U.S. Geological Survey currently recognizes the Lower-Upper Permian boundary within the Guadalupian Series at the Word-Capitan boundary, a decision not accepted now by most Permian workers.

The Road Canyon Formation was deposited during a time of maximum worldwide transgression of Permian seas; consequently, many units elsewhere in the world are correlated to it. In Texas, the Road Canyon contains the youngest of the Permian brachiopod bioherms (of Cooper and Grant, 1972, small moundlike organic buildups formed, in part, by brachiopods). A detailed biostratigraphic study was undertaken to resolve some of the apparent age anomalies. Because brachiopods, ammonoids, and fusulinids have produced differing age interpretations, an independent test was conducted by detailed sampling and study of conodonts from carefully measured sections.

THE TYPE SECTION (RC 1)

The Road Canyon Formation has been measured by several stratigraphers at several places (King, 1931, p. 71; Cooper and Grant, 1972, p. 67), and the sequence differs from place to place because of irregular carbonate buildups in the lower part of the unit (Cooper and Grant, 1972). The type section is located in the hill north of Leonard Mountain on the east side of Gilliland Canyon, due north of the Iron Mountain Ranch, slightly west of the 103°15' meridian (Cooper and Grant, 1964, fig. 1). This section, a part of King's (1931) Section 17, was measured in detail by John Cys and published in a guidebook (Cys, 1981). We followed Cys' section, in which the units are identified by painted numbers, taking 7-kg samples about every 10 ft

Manuscript approved for publication April 22, 1985

[1] U.S. Geological Survey, Reston, VA 22092
[2] Smithsonian Institution, Washington, DC 20560

Figure 1. Southwestern portion of the Glass Mountains showing section locations of the Road Canyon Formation.

where beds were exposed and clearly in place. Samples were not collected from silty zones, known from experience to be unlikely sources of conodonts. We then measured 11 more sections, east and west of the type section, to document the sequence of conodonts within the Road Canyon Formation.

Thickness measurements of the Road Canyon vary among workers, even when the identical section is measured. Major variations stem from differing choices of unit boundaries. There is little doubt about the lower boundary at the type section because the limestone of the Road Canyon occurs abruptly above the thick sequence of siltstone and mudstone of the Cathedral Mountain Formation. However, King (1931), Cooper and Grant (1964), Cys (1981), and the present writers all differ as to the upper boundary.

King (1931, p. 139, Section 17) defined the "First Limestone member" strictly on the ledge-forming limestone, 125 ft thick, at or near the type section of the Road Canyon. If significant higher limestone ledges that have intervening siltstone are included, the total thickness is about 208 ft. Cooper and Grant (1964, p. 1558) picked the upper boundary where siltstone becomes predominant, and limestone is present only as thin lenses and stringers and report a total thickness of 228 ft. By using the same criteria at the same section, we measured 232 ft, a small margin of error. Cys (1981, p. 200) picked the upper boundary at the last stringer of limestone and reported a total thickness of 282 ft; there is an error of 10 ft because the painted numbers show 245 followed by 260, rather than 250. Cys' measurement of 272 ft differs from ours only in our choice of the upper boundary; up to 232 ft, we replicated his measurements exactly. Our usage corresponds to the definition of the Road Canyon in its type section (Cooper and Grant, 1964) despite Cooper and Grant's statement that it was a direct equivalent of King's "First Limestone member" of the Word Formation.

Just above a basal conodont layer in the type section, a 50-ft-thick fossiliferous conglomeratic unit contains many brachiopods, chert nodules and lenses, algal sheets, and, especially, sponges. Rugose corals and "reef" or bank-forming brachiopods of the families Richthofeniidae and Lyttoniidae are abundant. Some beds in the upper part of the conglomeratic unit have aligned clusters of *Parathuramina*. The conodonts in this part of the unit are *Neogondolella idahoensis*, *Hindeodus excavatus*, and *Xaniognathus abstractus*. The conglomeratic unit is capped by 18 ft of packstone that has chert stringers and abundant crinoid debris; a sample taken at the 52.5-ft level contains the conodont species *Neogondolella idahoensis* and *N. serrata*. Above this level, *N. serrata* totally replaces *N. idahoensis*. Somewhat higher, above a bedded chert interval that Cys (1981, p. 200) reports capped by a local disaster, is a huge massive organic carbonate buildup of the kind that Cooper and Grant (1972) called a bioherm. This buildup is overlain by a 40-ft-thick yellow siliceous siltstone, from which no samples were taken. The top of the formation is placed at the carbonate ledge above which the thin-bedded siltstone typical of the Word Formation begins. Cys' (1981) selection of the uppermost of several carbonate lenses in the siltstone provides an ambiguous boundary for a distinct and mappable unit.

SUPPLEMENTARY SECTIONS

Section RC II is farther west on the same hill that contains the type section (fig. 1), where the slope faces westward into Gilliland Canyon and just above a small clay slide produced by erosion and slumping of the siltstone and mudstone of the Cathedral Mountain Formation. In Section RC II, the Road Canyon Formation is thinner than the type section—only 164 ft thick. As in the nearby type section, the formation begins with conglomeratic limestone that contains clasts of the Cathedral Mountain lithology and sponges, corals, bryozoans, and various brachiopods including such reef or bank-forming types as richthofeniids and the lyttoniid *Cancrinelloides*. *Cancrinelloides*, the "beaded" lyttoniid, is distinctive and characteristic of the Road Canyon, although it also occurs lower in the section in the Skinner Ranch and Cathedral Mountain Formations. The occurrence of this brachiopod supported the argument for a Leonardian age of the Road Canyon Formation (Cooper and Grant, 1972, 1973). In Section RC II, the top of the Road Canyon is placed at the base of a 43-ft interval of yellow-orange, thin-bedded siltstone, although thin lenses of pellodial packstone occur in the siltstone higher in the Word Formation.

One sample in Section RC II contains excellent examples of both *Neogondolella idahoensis* and *N. serrata*, thus revealing the point of transition between the two species. Samples below the 38-ft level contain only *N. idahoensis*, and those above that level contain only *N. serrata*. Beds immediately above and below that level are covered by soil; this cover prevents determination of the full extent of overlap in the ranges of the two species.

One other section (RC VII, 113 ft thick) was measured in the hill that contains the type section, but RC VII is much farther east on the Hess Ranch near the easternmost end of the hill (fig. 1). This section is near USNM (United States National Museum) locality 726c where large specimens of the ammonite *Perrinites*, a genus thought to be Leonardian in age (Miller and Furnish, 1957), were found by Cooper and Grant (1972, pl. 15, fig. 2). This occurrence of *Perrinites* provides one of the strong arguments for a Leonardian age of the Road Canyon Formation. The ostracode-form lyttoniid brachiopod *Coscinophora* occurs throughout the formation at this section and forms large banks in the lower part.

A thin bed below the lowest *Coscinophora* bank contains the conodont *Neogondolella idahoensis*, and a *Coscinophora* bed above the massive bank contains *N. serrata*; the change in conodont species is inferred to occur within the lower massive bank, about 35 ft above the base of the formation.

Farther east (fig. 1) on the Appel Ranch (now the Williams Bar-Crescent Ranch), the Road Canyon is only 6 ft thick (Sections RC VIII and RC XII), but the changeover in species of *Neogondolella* is contained within this thin unit. Section RC VIII is a conglomeratic limestone and forms a ledge along the north side of the canyon just north of the old Word Ranch site. Section RC XII is a limestone ledge east of Split Tank above USNM localities 726y, 726w, 702a', 702a, 702ent, 702un, 702low, 702inst, and 708a (Cooper and Grant, 1972, fig. 22). Transitional in thickness between these thin beds and the *Coscinophora*-rich beds of Section RC VII are Section RC X, located on the western side of hill 5611 near USNM locality 706, and Section RC XI, located on the knob that is USNM locality 702c.

Other sections are farther from the type section. Sections RC III and RC IV are in the hill, where a large carbonate buildup occurs, immediately west of Clay Slide (of King, 1931). Section RC IV is through the core of the carbonate mass, and Section RC III, which ends in a bedded chert-nodule dominated bank about 800 ft farther north. Only the conodont *Neogondolella serrata* is present throughout the formation at Sections RC III and RC IV. The change from *N. idahoensis* to *N. serrata* probably took place before carbonate deposition began; that is to say, in the siltstones and mudstones of the underlying Cathedral Mountain Formation.

Two sections were measured on the hill containing Clay Slide. Section RC V is on a subsidiary knob north of the slide itself, just above a smaller, grassier slump in the siltstones and mudstones of the Cathedral Mountain Formation. Both *N. idahoensis* and *N. serrata* were found in a sample taken 36.5 ft above the base of the formation. Section RC VI is immediately above Clay Slide, on the south flank of the hill. There *N. idahoensis* and *N. serrata* occur together at the 30-ft level, but a sample taken 50 ft above the base contained only *N. serrata* (along with species in other genera).

A very thick section (RC IX) was measured on the flank of Sullivan Peak. Here the Road Canyon is thin-bedded and silty and is apparently transitional to the deltaic facies that dominates farther to the northwest on Altuda (or Bird) Mountain (Davis and others, 1983). *N. idahoensis* was found at the base and about 90 ft above the base of the formation, and *N. serrata* and the brachiopod *Coscinophora* were found 246 ft above the base. The zone of transition between the two conodont species was not determined.

THE LEONARDIAN-GUADALUPIAN BOUNDARY

Species ranges of *Neogondolella* are the basis for conodont zonations of the Upper Permian and are important to zonation of the Lower Permian (Kozur, 1978). A relatively rapid changeover in species of *Neogondolella* occurs in the lower part of the Road Canyon Formation (this paper; Wardlaw and Grant, 1987). *N. idahoensis* is commonly found near the base, and *N. serrata* is found above the base. The species are separated by only a few feet or occur within the same sample in several sections, and the occurrences suggest that the changeover is rapid and overlap is small.

A similar relatively rapid changeover from *N. idahoensis* to forms identified as *N. serrata* was documented (Wardlaw and Collinson, 1984, 1986; Behnken and others, 1986) in the lower part of the Meade Peak Phosphatic Shale Member of the Phosphoria Formation in Wyoming and Idaho. Wardlaw and Collinson (1978, 1979) noted this change throughout the Great Basin Rocky Mountain region. Behnken (1975) showed a similar sharp change in faunas from the Guadalupe Mountains, Texas, where *N. idahoensis* occurs through most of the Victorio Peak and Bone Spring Limestones to the uppermost part of these units, and *N. serrata* occurs at the base of the overlying Cutoff Shale (Behnken, 1975, figs. 5, 6). All of the units that contain the *N. idahoensis* to *N. serrata* succession have been correlated by other fossil groups (for example, brachiopods, Cooper and Grant, 1972; Wardlaw and Collinson, 1978, 1986; ammonoids, Furnish, 1973; fusulinids, White and Todd, 1968) and are considered to be time correlatives. The biostratigraphic succession of *N. idahoensis* to *N. serrata* appears to be a widespread (at least over most of a continent) phenomenon that marks a nearly synchronous event.

We propose to define the Leonardian-Guadalupian boundary on the basis of the conodonts, the range of *N. idahoensis* is Leonardian, and the basal Guadalupian begins with the first occurrence of *N. serrata*. The Road Canyon Formation spans the boundary, and the concept of a Roadian Stage ceases to be useful. Consequently, we advocate abandonment of Roadian as a stage. This new boundary definition also conforms more closely to the original definition of the Leonard and Word Formations, upon which the stages are based, by including most of the Road Canyon (First Limestone member of the Word) in the Wordian (basal Guadalupian).

Our conclusions are twofold. (1) The Leonardian-Guadalupian boundary can be defined as the level at which *N. idahoensis* is replaced by *N. serrata*. (2) The concept of a Roadian Stage should be abandoned.

DEPOSITIONAL SETTING

King (1942, fig. 18) (fig. 2) presented the generally accepted Permian regional geography of West Texas. The Permian sediments of the Glass Mountains were deposited on the narrow southern shelf between the Marathon folded belt, which formed a land area to the south, and the Hovey channel and Delaware basin, which formed a relatively deep marine channel and basin to the north. King (1931), in his classic study of the Glass Mountains, noted that the distribution of sediments in Leonard and Word time showed that sandstone, conglomerate, and redbeds were thickest near the uplifted area to the south and thinned to the north and northeast where they were replaced laterally by limestone (King, 1931, p. 89). This configuration is one of inner shelf clastics and outer shelf carbonates deposited on a very narrow shelf.

Figure 2. Paleogeographic setting of the Permian of West Texas (from King, 1942) showing location of the Glass and northern Del Norte Mountains.

Cys (1981) differed with this interpretation radically by postulating a shelf boundary perpendicular to that of King's (Cys, 1981, fig. 11; Cys and Mazzullo, 1978, fig. 5) and by interpreting the Leonard sediments as being largely slope and basin deposits. Tabulations indicating basinal deposition and thin limestones to the northeast indicating shallow shelf deposition.

Cooper and Grant (1972) agreed with King's general shelf deposition for the Leonard and Word sediments. Further, they noted that, to the southwest in the Del Norte Mountains (fig. 2) in late Cathedral Mountain time, local uplift shed coarse conglomerates composed of well-rounded pebbles. Abundant transported ammonoids, fragmentary brachiopods, and numerous plant stem fragments indicated a shore zone (Cooper and Grant, 1972, p. 99). Rohr and others (1982) and Davis and others (1983) interpreted the conglomerates and related siltstones in the Del Norte Mountains to be nonmarine fluvial and marine deltaic deposits. Many delicate, well-preserved leaves are present in the siltstones (Murray and others, 1984; Rohr and others, 1987), and these sediments are equivalent to the upper part of the Cathedral Mountain and Road Canyon Formations (Murray and others, 1984; Wardlaw, unpub. conodont data).

The Road Canyon Formation is, perhaps, the most poorly known unit in the Glass Mountains. Although Cys (1981) did not interpret the depositional environment for the Road Canyon Formation in his Leonardian study, he showed it unconformably overlying "basinal" deposits of the Cathedral Mountain Formation.

The Road Canyon Formation represents widespread carbonate deposition during transgression apparently related to a worldwide sea-level rise (Wardlaw, 1980). The Road Canyon interfingers with the deltaic and fluvial deposits of the Del Norte Mountains and probably represents relatively shallow-water deposition. In many places, the Road Canyon is conglomeratic both at its base and throughout much of its thickness. Clasts include irregular limestone and siltstone cobbles that are either penecontemporaneous or are derived from the underlying Cathedral Mountain Formation and flat siltstone cobbles that are derived from the underlying Cathedral Mountain Formation. In no instance do the limestone clasts and matrix yield different conodont faunas. The clasts indicate active reworking of underlying and penecontemporaneously lithified sediments and very little transport; we interpret this to be indicative of shallow-water deposition.

Carbonate buildups within the Permian of the Glass Mountains have been interpreted as being either in place or redeposited (see Cys and Mazzullo, 1978, and Flores-McMillan and Watters, 1978). Sedimentologists with one exception (or another) seem to select buildups that tend to support their own interpretations. Each buildup should be analyzed separately; many are in place and many are not. The *Cuifloquina* bank at RC IV, the flank deposits at RC III, and the *Cuifloquina* banks at RC VII are in-place buildups. The more diversified buildups at the base of RC I and RC IX may be redeposited. In spite of a diversity of opinions, we feel that the depositional features indicate shallow-water deposition.

CONODONT BIOFACIES

Wardlaw and Collinson (1984) showed conodont biofacies distributions for the Permian Phosphoria Formation and related rocks in the Rocky Mountains. The Road Canyon Formation and the lower part of the Phosphoria Formation were deposited during the same time interval (see Behnken and others, 1986), and both formations contain the same conodont species. Generally, Wardlaw and Collinson found that *Hindeodus* and *Sweetina* characterize nearshore environments and *Neogondolella* and *Xaniognathus* characterize offshore environments. Following the distribution model of Wardlaw and Collinson (1984), the Road Canyon Formation would have been deposited in intermediate to offshore environments because there is a mixing of offshore and nearshore elements. *Neogondolella* and the complete seximembrate apparatus of *Xaniognathus* are better represented than the seximembrate apparatuses of *Hindeodus* and *Sweetina*, probably due to transportation of the latter. The distribution of conodonts supports offshore deposition.

Neostreptognathodus, which characterizes intermediate environments in the Phosphoria Formation, is very rare in the Road Canyon Formation. *Sweetognathus*, which is very rare in the Phosphoria, is also rare in the Road Canyon but is more common than *Neostreptognathodus*. Perhaps these differences can be explained by the narrow depositional shelf that must have existed during the deposition of the Road Canyon as compared to the broad depositional shelf that existed during the deposition of the Phosphoria. The narrow shelf would have compressed the biofacies, made differentiation of the biofacies less clear, increased overlap of the biofacies, and perhaps been unfavorable for such intermediate forms as *Neostreptognathodus*. Offshore shelf deposition for the Road Canyon Formation fits well with the original depositional interpretations of King (1931).

The ratios of number of samples containing *Hindeodus* or *Sweetina* or both to number of samples containing *Neogondolella* or *Xaniognathus* or both in each section (table 1) show that *Neogondolella* and *Xaniognathus* are generally more common than *Hindeodus* and *Sweetina*. Section RC IV is the exception, because *Hindeodus* and *Sweetina* are more common than *Neogondolella* and *Xaniognathus*, possibly indicating very shallow-water deposition. Also shown in table 1 is the percentage of samples that yielded conodonts in each section. Samples were generally of equal size and averaged 7 kg. Apparently, two factors severely limit the recoverability of conodonts. Sections RC

Table 1. HOS:NOS ratio and percent of samples that yielded conodonts from sections of the Road Canyon Formation.

[HOS, number of samples containing *Hindeodus* or *Sweetina* or both; NOS, number of samples containing *Neogondolella* or *Xaniognathus* or both]

Section	Yield, in percent	HOS:NOS
RC I	88	0:64
RC II	90	5:0
RC III	91	8:6
RC IV	43	1:50
RC V	50	5:0
RC VI	90	6:7
RC VII	67	6:7
RC VIII	80	1:10
RC IX	50	3:3
RC X	100	1:10
RC XI	100	1:10
RC XII	100	1:10

V, RC VI, and RC IX (figs. 1, 3) are dominated by platy and thin-bedded, silty lime-mudstone and siltstones and have poor conodont recovery, reflecting a greater influx of silt during deposition. Sections RC IV and RC VII (figs. 3, 4) are primarily *Coscinophora*-dominated carbonate buildups that have poor conodont recovery, probably reflecting very shallow water deposition.

SYSTEMATICS

Upper Paleozoic conodont species have been shown by Merrill and Powell (1980) to be variably represented by apparatuses. Although most individuals apparently start with a full complement of elements (generally six different kinds), some lose elements during ontogeny. Ontogenetic loss appears to vary among species and between environments but seems most variable between genera. For all practical purposes, genera can be characterized as (1) those that retain a full complement of elements through life, (2) those that lose a few elements (rare), (3) those that commonly lose most of the elements but show rare cases of total retention, and (4) those that occur only as a single element and do not commonly have a full complement of elements. In our material, *Hindeodus*, *Sweetina*, and *Xaniognathus* appear to be represented by seximembrate apparatuses (type 1). *Neostreptognathodus* and *Sweetognathus* appear to be represented by rarely occurring seximembrate apparatuses; both are generally represented by just the Pa element (type 3). *Neogondolella* appears to be represented by a single element platform (Pa) apparatus that probably occurred as a pair (type 4). The occurrence of conodont species in our material is shown in table 2.

Genus *Hindeodus* Rexroad and Furnish, 1964

Type Species.—*Hindeodus cristulus* (Youngquist and Miller, 1949).

Rexroad and Furnish, 1964, p. 671–672.

Hindeodus excavatus (Behnken)

Plate 2, figures 1–15; plate 3, figures 3–11.

Ellisonia excavata Behnken, 1975, p. 302, pl. 1, figs. 9–14.

Anchignathodus minutus (Ellison) Behnken, 1975, p. 297, pl. 1, figs. 16–18.

Hindeodus excavatus (Behnken); Sweet, 1977a, p. 215, pl. 1, figs. 7–11; Wardlaw and Collinson, 1984, p. 268–269, pl. 5, figs. 1, 2, 4–9; 1986, p. 133, fig. 17.13–20.

Diagnosis.—A species of *Hindeodus* characterized by a Pa element with a large cusp; denticles increase in width anteriorly, except for the anteriormost, and generally decrease in height anteriorly, except for the posteriormost three, which may be of subequal height; cusp is much higher than denticles; and an Sa element with short lateral processes that are slightly upturned laterally and that bear for at least part of their length denticles of alternating (smaller versus larger) sizes (see Behnken, 1975, p. 299, pl. 1, fig. 12). The entire apparatus is illustrated in Sweet (1977a) and Wardlaw and Collinson (1984, 1986).

Discussion.—*H. excavatus* is commonly represented in our collections by the Pa element and a few broken ramiform elements. Of these, Sc elements are most common, followed by M, Sb, and Pb elements. No Sa element was recovered. We interpret the incomplete and broken condition of the elements of *Hindeodus* to indicate that the skeletal remains of this species were probably deposited in high-energy environments and, in part, were probably transported short distances. *H. excavatus* appears to be the only conodont common to the *Coscinophora* banks that we also interpret as being deposited in high-energy and very shallow water environments.

Genus *Neogondolella* Bender and Stoppel, 1965

Type Species.—*Neogondolella mombergensis* (Tatge), 1956, pl. 6, figs. 1, 2.

Neogondolella idahoensis (Youngquist, Hawley, and Miller)

Plate 1, figures 8, 9, 14–16, 19–22; plate 2, figures 20–22, 26–28; plate 4, figures 24, 25.

Gondolella idahoensis Youngquist, Hawley, and Miller, 1951, p. 261, pl. 54, figs. 1–3, 14, 15; Clark and Ethington, 1962, p. 108, pl. 2, figs. 15, 16; Clark and Mosher, 1966, p. 388, pl. 47, figs. 9–12; Clark and Behnken, 1971, p. 431, fig. 9; Kozur, 1978, pl. 5, figs. 2, 15, 16.

Neogondolella idahoensis (Youngquist, Hawley, and Miller); Behnken, 1975, p. 306–307, pl. 1, figs. 28–30; Wang, 1978, p. 220–221, pl. 2, figs. 25, 26; Igo, 1981, p. 38, pl. 1, figs. 11–13, 15, 16; Wardlaw and Collinson, 1984, p. 269, pl. 1, figs. 10, 11, 1986, p. 133, fig. 17.11, 12; Behnken, Wardlaw, and Stout, 1986, p. 179–181, fig. 4.12–16, 18.

Gondolella phosphoriensis Youngquist, Hawley, and Miller (part), 1951, p. 262, pl. 54, figs. 27, 28; Clark and Ethington, 1962, p. 108, pl. 2, figs. 17–18.

Diagnosis.—A species of *Neogondolella* characterized by a platform of moderate width that gradually narrows in the anterior two-thirds of its length and has a blunt, squared posterior platform margin; denticles that generally increase in size anteriorly, except for the anterior one or two; and well-developed furrows.

Discussion.—Our examples of *N. idahoensis* appear to be very similar to those previously described from the Phosphoria Formation in Idaho (Youngquist, Hawley, and Miller, 1951) and the Bone Spring Limestone in West Texas (Clark and Ethington, 1962; Behnken, 1975). This species seems reasonably consistent in morphology throughout its worldwide distribution.

Neogondolella serrata (Clark and Ethington)

Plate 1, figures 1–7, 10, 13, 17, 18; plate 2, figures 16–19, 23–25.

Gondolella serrata Clark and Ethington, 1962, p. 108–109, pl. 1, figs. 10, 11, 15, 19; pl. 2, figs. 1, 5, 8, 9, 11–14; Clark and Mosher, 1966, p. 389, pl. 47, figs. 13–15; Clark and Behnken, 1971, p. 431, pl. 1, fig. 10; Kozur, 1978, pl. 5, fig. 22.

Neogondolella serrata (Clark and Ethington); Wang, 1978, p. 222, pl. 2, figs. 6–8, 14, 15, 20–22; Clark and Behnken, 1979, p. 268–271, pl. 1, fig. 12; Wardlaw and Collinson, 1984, p. 270, pl. 2, figs. 13–15, 1986, p. 133–134, fig. 17.7, 8; Behnken, Wardlaw, and Stout, 1986, p. 183, fig. 5.20, 21, 23–31.

Neogondolella serrata serrata (Clark and Ethington); Behnken, 1975, p. 308, pl. 2, figs. 21–24, 37.

Diagnosis.—A species of *Neogondolella* characterized by a platform of moderate width, which is generally short and arched, narrows anterior to the middle, and has a round to blunt posterior platform margin; lateral platform margin variably serrated, commonly with at least three or more paired serrations on the anterior, tapering part of platform; rarely serrated for almost entire margin; a moderate cusp of circular cross section; denticles generally increasing in size anteriorly except commonly one or two smaller denticles along middle of carina and one or two denticles at anterior end; and moderately developed furrows.

Discussion.—*N. serrata* appears to show the earliest regional morphologic variation that begins provincialism in Permian neogondolellids. Specimens from Idaho can be distinguished from those of West Texas, although in both regions *N. serrata* appears to replace *N. idahoensis* in remarkable synchroneity. The younger species of *Neogondolella* in Idaho are completely different from those of West Texas, and so *N. serrata* appears to be the youngest species to show worldwide distribution; however, it presages the later trend of provincialism by developing distinct geographic morphotypes.

Genus *Neostreptognathodus* Clark, 1972

Type Species.—*Neostreptognathodus sulcoplicatus* (Youngquist, Hawley, and Miller, 1951).

Clark, 1972, p. 153.

Neostreptognathodus clinei Behnken

Plate 3, figure 2; plate 4, figure 23.

Neostreptognathodus clinei Behnken, 1975, p. 309–310, pl. 2, figs. 15, 16; Sweet, 1977b, p. 237, pl. 1, fig. 4; Wardlaw and Collinson, 1984, p. 270, pl. 5, fig. 13; Behnken, Wardlaw, and Stout, 1986, p. 183, fig. 4.6, 7.

Diagnosis.—A species of *Neostreptognathodus* characterized by a platform element (Pa) with two carinae that are smooth or only faintly ornamented posteriorly by poorly developed nodes; carinae separated by a distinct groove; blade joins platform at center.

Discussion.—*N. clinei* is a rare long-ranging form that is found throughout North America in upper Leonardian and lower Guadalupian strata. A Pb element recovered from the lower part of the Word Formation, Section RC I, belongs to the apparatus of *Neostreptognathodus* (see Wardlaw and Collinson, 1986). It is tentatively assigned to *N. clinei* and illustrated (pl. 4, fig. 23) to compare with Pb elements of *Xaniognathus*.

Neostreptognathodus newelli Wardlaw and Collinson

Plate 3, figure 1.

Neostreptognathodus newelli Wardlaw and Collinson, 1984, p. 270, pl. 5, figs. 10–12, 14–17; 1986, p. 135, fig. 18.4.

Neostreptognathodus sp. C, Baird, 1975, p. 41–42, pl. 1, figs. 11, 12; Wardlaw and Collinson, 1978, p. 1177, 1979, p. 152.

Diagnosis.—A species of *Neostreptognathodus* characterized by a platform element (Pa) with two carinae ornamented by 12 to 17 pairs of slightly oblique, low transverse ridges that are separated by a narrow but distinct median groove; posteriormost one to three ridges merge, terminating the median groove; blade joins platform slightly off center (in contrast to *N. sulcoplicatus* and *N. clinei*, which are centered) and has a poorly developed groove, which separates the blade from the closest carina.

Discussion.—*N. newelli* is rare in the Road Canyon Formation and occurs only in one sample. It is common throughout most of the Great Basin and northern Rocky Mountains and is diagnostic of the *Penicuiaurris basis-Neostreptognathodus newelli* Zone (formerly *N.* sp. C) of Wardlaw and Collinson (1978) that they correlated with the upper part of the Road Canyon. This fortuitous recovery strengthens that correlation. The *Penicuiaurris basis-Neostreptognathodus newelli* Zone is the near-shore time equivalent of the *Neogondolella serrata* Zone, found in the Phosphoria Formation in Idaho and in the Cutoff Shale and Road Canyon Formation in West Texas.

Genus *Sweetina* Wardlaw and Collinson, 1986

Type Species.—*Sweetina triticum* Wardlaw and Collinson, 1986, p. 135–138.

Sweetina festiva (Bender and Stoppel)

Plate 3, figures 18–25.

Lonchodina festiva Bender and Stoppel, 1965, p. 345–346, pl. 15, figs. 9, 10.

Stepanovites festivus (Bender and Stoppel); Kozur, 1975, p. 23; Kozur, 1978, pl. 5, fig. 29; Kozur and Mostler

isbech, 1979, p. 119, pl. 4, figs. 1, 2; Wardlaw and Collinson, 1984, p. 271, pl. 3, figs. 4–11).

Xaniognathus fertilis (Bender and Stoppel); Wardlaw and Collinson, 1986, p. 138.

Diagnosis.—A species of *Xaniognathus* characterized by a pastinate Pa element with a poorly developed lateral process and a dygytate M element with a very large cusp, one to three anterolateral denticles, generally no posterolateral denticles, and a greatly flared triangular basal cavity that is directed posterolaterally.

Discussion.—The entire apparatus of *Xaniognathus triticum* is illustrated in Wardlaw and Collinson (1986, fig. 20). *X. fertilis* appears to have an apparatus that is very similar to that of *X. triticum* except for the Pa element. The Pa element of *X. fertilis* has a longer lateral process, a less prominent cusp, and a less well developed basal cavity than the Pa element of *X. triticum*. *X. fertilis* is commonly represented in our material by broken ramiform and platform elements. Sc elements are the most common followed by fairly common M elements, and then by Pa, Sb, and Sa elements. No Pb elements were recovered. The occurrence of *X. fertilis* is thought to be similar to that of *Hindeodus excavatus*.

Genus *Sweetognathus* Clark, 1972

Type Species.—*Sweetognathus whitei* (Rhoades, 1963); Clark, 1972, p. 155.

Sweetognathus iranicus Kozur, Mostler, and Rahimi-Yazd

Plate 5, figures 12–17

Sweetognathus iranicus Kozur, Mostler, and Rahimi-Yazd, 1975, p. 9–10, pl. 4, figs. 1–10, pl. 5, figs. 2–4; Kozur, 1978, pl. 6, figs. 4, 7–10, 11, 15, 18.

Diagnosis.—A species of *Sweetognathus* characterized by a Pa element with rounded denticles or nodes that have postulose microornament forming a row of small nodes connected by a thin postulose longitudinal ridge that is laterally, not centrally, placed and a variably developed secondary row of nodes that generally through growth become larger than, and merge with, the primary row to form transverse ridges.

Discussion.—*S. iranicus* is rare in the Road Canyon Formation. A partial apparatus has been illustrated for *Sweetognathus* (Ritter, 1986). Our sparse material appears to be represented only by Pa elements. *S. iranicus* was originally described from the Abadeh section of central Iran (Kozur, Mostler, and Rahimi-Yazd, 1975). *S. iranicus* appears to range throughout the Road Canyon Formation and into the lower part of the Word Formation in West Texas.

Genus *Xaniognathus* Sweet, 1970

Type Species.—*Xaniognathus curvatus* Sweet, 1970, p. 262.

Xaniognathus abstractus (Clark and Ethington)

Plate 4, figures 1–22

Subbryantodus abstractus Clark and Ethington, 1962, p. 112–113, pl. 1, figs. 16, 20, 21, pl. 2, fig. 2.

Lonchodina mulleri Tatge; Clark and Ethington, 1962, p. 110–111, pl. 1, fig. 4.

Apatognathus tribulosus Clark and Ethington, 1962, p. 107, pl. 1, figs. 3, 7, 13, 17.

Xaniognathus abstractus (Clark and Ethington); Behnken, 1975, p. 313, pl. 1, fig. 15; Wardlaw and Collinson, 1984, p. 271, pl. 2, figs. 1–12; Wardlaw and Collinson, 1986, p. 135, fig. 18.17–23; Behnken, Wardlaw, and Stout, 1986, p. 185, figs. 8.4–7, 9.7–20.

Xaniognathus sp. Malkowski and Szaniawski, 1976, p. 83, pl. 1, fig. 2.

Xaniognathus sentis (Tatge); Behnken, 1975, p. 313, pl. 2, fig. 13.

Xaniognathus curvatus Szaniawski, 1976, p. 254, pl. 9, figs. 1–7.

Ozarkodina fertilis Tatge; Clark and Behnken, 1971, p. 433, pl. 2, fig. 8.

[A complete element synonymy is given in Behnken, Wardlaw, and Stout, 1986]

Diagnosis.—A species of *Xaniognathus* characterized by the delicate, greatly recurved Sb (*Hindeodus rebuliosus*) element with large cusp and low denticles on one process and a large variable Pb element with a twisted posterior process commonly bearing five to eight denticles; rare specimens have more denticles, and an anterior process commonly bearing 7 to 11 or more denticles, with midlateral rib for most of its length; anterior process variable, robust to relatively thin and delicate. The entire apparatus is illustrated in Wardlaw and Collinson (1984, pl. 2, 1986, fig. 18) and Behnken, Wardlaw, and Stout (1986, fig. 8).

Discussion.—*X. abstractus* is abundant and well preserved in many of our samples. Sb, M, and Pa elements are the most commonly recognized, although many undetermined fragments probably represent Sc elements. Sa and Pb elements are slightly less common. The complete representation and excellent preservation of *Xaniognathus* indicate that it is a truly indigenous faunal element.

Table 2. Occurrence of conodont species in samples of the Road Canyon Formation sections that include underlying and overlying units.

[1, *Anagnostiella lapidosa*; 2, *Neogondolella serrata*; 3, *Hindeodus excavatus*; 4, *Sweetognathus iranicus*; 5, *Xaniognathus fertilis*; 6, *Xaniognathus abstractus*; 7, *Xaniognathus curvatus*; 8, *Neogondolella rosenkrantzi*; 9, *Neogondolella subcarinata*]

Sample	Lithology	Conodonts								
		1	2	3	4	5	6	7	8	9
	RC 1									
WS1181	0.5	X			X		X			
WS1182	7.5	X			X	X				
WS1183	26.0				X		X			
WS1184	40.0									
WS1185	52.4	X	X	X	X	X				
WS1186	68.0		X		X					
WS1187	77.5					X				
WS1188	88.0		X		X		X			
WS1189	100.0		X							
WS1170	155.0		X				X			
WS1171	170.0		X				X			
WS1172	192.0		X				X			
WS1173	212.0						X			
WS1174	229.0		X	X			X			
WS1175	250.0		X	X			X			
WS1176	272.0		X	X	X	X	X	X		
	RC II									
WS1179	61.0	X		X	X		X			
WS1180	0			X						
WS1188	12.0									
WS1187	38.0	X	X				X			
WS1186	61.0		X		X					
WS1185	109.0		X				X			
WS1184	129.0									
WS1183	144.0		X							
WS1182	184.0		X	X						
WS1181	211.0		X	X	X					
WS1180	234.0		X				X			
	RC III									
WS1198	0						X			
WS239	2.8					X				
WS240	4.0		X				X			
WS1192	17.0			X						
WS1193	28.0						X			
WS1194	52.0		X	X						
WS1195	59.0			X						
WS1196	45.0		X	X	X		X			
WS1197	58.0		X	X			X			
WS1198	74.0						X			
WS1199	88.0									
	RC IV									
WS1200	2.5		X	X		X				
WS241	5.0		X	X		X				
WS1201	17.0									
WS1202	39.0									
WS1203	67.0			X						
WS1204	80.0									
WS1205	98.0									
	RC V									
WS1209	4.8			X			X			
WS1208	16.5	X	X							
WS1207	43.0									
WS1206	77.0									
	RC VI									
WS1214	5.0									
WS1215	40.0	X	X	X						

Table 2. Occurrence of conodont species in samples of the Road Canyon Formation sections that include underlying and overlying units—Continued

[1, Neostreptognathodus sulconiger; 2, Neogondolella aserrata; 3, Hindeodus excelsus; 4, Sweetina hollina; 5, Neostreptognathodus clarkei; 6, Xaniognathus altanicus; 7, Neostreptognathodus clinei; 8, Neostreptognathodus nevadi; 9, Neostreptognathodus sulconiger]

Sample	Footage	Conodonts								
		1	2	3	4	5	6	7	8	9
W81232	80.5	—	X	—	—	—	X	—	—	—
W81231	72.0	—	—	—	—	—	—	—	—	—
W81230	67.0	—	—	X	—	—	X	—	—	—
RC VII										
W81217	1.5	X	—	X	X	—	X	—	—	—
W81218	13.5	X	—	—	—	—	X	—	—	—
W81219	18.0	—	—	—	—	—	—	—	—	—
W827	19.0	—	—	—	—	—	—	—	—	—
W828	27.0	—	—	X	—	—	—	—	—	—
W81220	35.0	—	—	—	—	—	—	—	—	—
W829	35.0	—	—	X	—	—	—	—	—	—
W81221	49.0	—	—	—	—	—	—	—	—	—
W81222	70.0	—	X	—	—	—	—	—	—	—
W81223	89.0	—	X	—	—	—	—	—	—	—
W81224	112.0	—	X	—	—	—	—	—	—	—
W81215	130.0	—	X	X	X	—	X	—	—	—
RC VIII										
W81226	1.5	—	—	X	X	—	—	—	—	—
W81227	26.0	X	—	X	X	—	X	—	—	—
W8210	27.5	—	—	X	—	—	X	—	—	—
W8211	28.5	—	—	—	—	—	—	—	—	—
W8212	29.5	X	—	—	X	—	X	—	—	—
W81228	30.5	—	X	X	X	—	X	—	X	—
W81229	39.0	—	—	—	—	—	—	—	—	—
W81230	59.0	—	—	—	—	—	—	—	—	—
RC IX										
W81244	0.0	X	—	—	—	—	X	—	—	—
W81243	5.0	X	—	—	—	—	X	X	—	—
W81241	74.0	X	—	—	—	—	—	—	—	—
W8242	95.0	—	—	—	—	—	—	—	—	—
W8244	114.0	—	—	—	—	—	—	—	—	—
W81242	144.0	—	—	—	—	—	—	—	—	—
W8245	160.0	—	—	—	—	—	—	—	—	—
W8246	195.0	—	—	—	—	—	—	—	—	—
W81236	246.0	—	X	X	—	—	X	—	—	—
W81237	251.5	—	X	X	—	—	—	—	—	—
W81238	272.5	—	X	—	—	—	—	—	—	—
W81239	281.5	—	X	X	—	—	X	—	—	—
W81240	291.0	—	X	—	—	—	X	—	—	—
RC X										
W877	1.0	—	X	X	X	—	X	—	—	—
W878	29.0	—	X	X	—	—	—	—	—	—
W879	52.0	—	X	X	—	—	X	—	—	—
RC XI										
W81225	0.5	X	X	X	X	—	X	—	X	X
W876	5.5	—	X	X	X	X	X	—	—	—
702C	5.5	—	X	X	—	—	X	—	—	—
RC XII										
W8471	0.5	X	—	X	X	X	X	—	—	X
W8472	5.5	—	X	X	X	X	—	—	—	—

[Four specimens of Neostreptognathodus sulconiger]

MEASURED SECTIONS

Section RC I (fig. 4)

Type section of the Road Canyon Formation. Hill north of Leonard Mountain, east side of Gilliland Canyon, west of 103°15′ meridian. Compare with sections by Cooper and Grant (1964) and Cys (1981) at the same locality.

	Thickness (ft)	Conodont sample
Word Formation:		
11. Siltstone, yellow, vitreous, with thin- to medium-bedded limestone stringers; conodont sample at 272 ft	28	272
10. Siltstone, yellow, vitreous, capped by 4-ft bed of skeletal packstone with abundant brachiopod spines, brachiopods, echinoids, ammonoids, corals, bryozoans, sponges; original limestone; conodont sample at 250 ft	26	252
Road Canyon Formation:		
9. Limestone, gray, rubbly weathering, medium- to thick-bedded, silty packstone with some chert; most pebbles are algal coated; algal fenestrals, corals, snails, brachiopods (*Linoproductus* at 230 ft); conodont samples at 226, 212, 192 ft	42	232
8. Siltstone, yellow, vitreous, thin-bedded, thin lenses of medium-brownish-gray, silty wackestone with *Pseudoschwagerina*; conodont sample at 170 ft	23	190
7. Limestone, ledge, medium-brownish-gray skeletal packstone with fossil bed at 164 ft containing brachiopods (*Cleiothyridina*, *Derbyia*, *Neospirifer*), bryozoans, *Meekella*, sponges, corals, echinoid debris; sponge spicules; conodont sample at 155 ft	33	167
6. Siltstone, yellow, vitreous, shaly-bedded with thin beds of pelsoid packstone	27	134
5. Siltstone with thin beds of pelsoid packstone changing upward to packstone with thin stringers of siltstone; fenestrate, crinoid columnals; conodont sample at 110 ft	46	127
4. Limestone, massive, organic carbonate buildup; medium-brownish-gray wackestone; sponges, corals, bryozoans, brachiopods (*Composita*, *Cleiothyridina*, *Derbyia*, *Meekella*, *Neospirifer*); conodont sample at 86 ft; barren sample at 77.5 ft	32	87
3. Chert, bedded, interbedded by overlying carbonate, medium-brownish-gray, weathers light brown, spicules; top is Cys "basal datum"	9	78
2. Limestone, brownish-medium-gray packstone with rusty chert stringers of skeletal fossil, medium-bedded; conodont samples at 69, 52.5 ft	18	69
1. Conglomerate, limestone, medium-brownish-gray, weathers orange to brown, packstone to wackestone; chert as boulders and cobbles, clasts from pebble to boulder size; limestone and chert fossils mostly sponges and corals, also crinoid columnals, brachiopod (*Linoproductus*), *Cordaites*	51	51

Section RC II (fig. 5)

Measurements made on the southwest-facing side of Hill 6779, the same hill that contains the type section on its south side; photo on eroded scar of Cathedral Mountain siltstone and mudstone that forms a small clay slide.

	Thickness (ft)	Conodont sample
Word Formation:		
9. Siltstone, yellow-orange, thin-bedded, very localized lenses and beds of pelsoid packstone up to 4 ft thick appear to be channel deposits; corals and other fossils; conodont sample at 254–233 ft	57	244
8. Siltstone, yellow-orange, thin-bedded, minor amounts of limestone, partly covered	43	207
Road Canyon Formation:		
7. Limestone, light-brownish-gray, coarse-grained pelsoid packstone with abundant colonial debris; few brachiopods (*Meekella*, *Orthotichia*); a massive carbonate buildup that becomes bedded and silty toward top; conodont samples at 164, 144, 129, 109 ft	80	164
6. Covered, probably thin-bedded siltstone	40	104
5. Limestone, medium-brownish-gray pelsoid skeletal packstone with rusty brown silica-fied "skins" on bedding planes and joints; conodont sample at 61 ft	17	64
4. Covered, probably shaly siltstone	8	47
3. Limestone, medium-olive-gray, fine-grained skeletal packstone with corals, bivalves, crinoid columnals, echinoderm fragments, brachiopods (*Composita*, *Cleiothyridina*, *Derbyia*, *Dictyoclostus*, *Meekella*, *Neospirifer*, *Perditocardinia*); nodular, base to 0.5 ft of medium-orange-pink crinoidal packstone; conodont samples at 26, 7.5, 0.5 ft; barren sample at 40 ft	50	50

Section RC III (fig. 5)

Hill west of hill containing Clay Slide; bedded flank deposits about 400 ft north of crest of massive carbonate buildup

	Thickness (ft)	Conodont sample
Road Canyon Formation:		
8. Limestone, medium-gray, sandy, skeletal lime-		

Figure 3. Columnar sections of the Road Canyon Formation. Sections RC IX, RC IV, RC III, RC VI, RC V, and RC II. Lithic symbols as in figure 4.

Figure 4. Columnar sections of the Road Canyon Formation. Sections RC I, RC VII, RC X, RC XI, RC VIII, and RC XII. RC I is the type section of the Road Canyon Formation. Datum is the sharp changeover from *Neogondolella idahoensis* to *N. serrata*.

mudstone to wackestone with chert nodules and beds; abundant corals, bryozoans, crinoid debris, fenestrate, spinose, and thick-shelled brachiopods (*Derbyia*, *Meekella*); conodont samples at 86, 74, 60 ft. | 54 | 40
7. Limestone, medium-gray, fine mudstone with chert lenses, medium-bedded, partly covered. | 10 | 36
6. Limestone, medium-gray, fine mudstone, medium bedded with brachiopods (*Neospirifer*, abundant *Meekella*); conodont sample at 45 ft. | 6.5 | 46
5. Limestone, orange-brown-gray, silty skeletal wackestone to packstone, bryozoans, crinoid, colonnade, brachiopods (*Rhombella*); chert nodules (not bedded) represents a thin finger of the carbonate buildup into Clay Slide deposits; conodont sample at 39 ft. | 1.5 | 39.5
4. Limestone, medium-gray, sandy, lime mudstone, fine grained corals, clams, bryozoans, brachiopods (*Neospirifer*, abundant *Meekella* at 37 ft); conodont sample at 32 ft, barren sample at 28 ft. | 10 | 38
3. Limestone, orange-gray skeletal wackestone, thick-bedded with rusty colored, sponges, carbonate pebbles and cobbles, corals, sponges, brachiopods (*Cleiothyridina*, *Derbyia*); represents a finger of the carbonate buildup into flank deposits; conodont sample at 17 ft. | 2.5 | 28
2. Covered, probably sandy, silty carbonate. | 11 | 18
1. Limestone, medium-grey lime mudstone with chert lenses, fine sand, scattered corals, brachiopods; conodont samples at 4, 2.5, 1 ft. | 4 | 4

Section RC IV (Fig. 1)

Hill west of Clay Slide, through massive carbonate buildup about 400 ft south of Section RC III.

	Thickness (ft)	Curve letter
Road Canyon Formation		
6. Conglomeratic limestone, orange-gray wackestone, massive bed with sponges, *Cleiothyridina*; barren conodont sample at 98 ft.	4	104
5. Partly covered, silty conglomeratic wackestone.	11	95
4. Limestone, orange-gray wackestone, *Cleiothyridina* bank, and some conglomeratic barren; conodont sample at 80 ft.	5	84
3. Limestone, light gray to orange, silty, crinoidal, dolomitic wackestone with chert nodules, massive, becoming conglomeratic upward; *Cleiothyridina* bank at 65 ft; conodont sample at 67 ft.	30	80
2. Dolostone, light-brownish-gray, silty, crinoidal dolowackestone, massive, localized lens with rusty chert nodules.	11	50
1. Limestone, light-olive-gray, skeletal wackestone to packstone, crinoidaceous, locally conglomeratic with clasts within clasts; sponges, corals, bryozoans, *Cleiothyridina* banks at 33, 25, 15 ft; conodont samples at 3, 2.5 ft; barren samples at 39, 17 ft.	39	39

Section RC V (Fig. 2)

Knoll hill that connects Clay Slide, but on knob north of the main, unnamed Cathedral Mountain siltstone also mudstone capped by Road Canyon limestone.

	Thickness (ft)	Curve letter
Road Canyon Formation	32	32
5. Limestone, silty lime mudstone with a general rusty chert nodules, thin, platy bedded, few fossils, corals, crinoid columnals, brachiopods, forming large capping hill; barren conodont samples at 77, 49 ft.	11	43
4. Limestone, cherty, silty lime mudstone, platy and wavy bedded, chert in many layers with sparse fine materials of silty skeletal wackestone with crinoid debris and brachiopods; conodont sample at 46.5 ft.	16	48
3. Chert and limestone, interbedded, tan, very fine grained, sandy, lime mudstone, and chert, chert probably replaced other layers, thin wavy bedded chert dominating upward.	24	30
2. Conglomeratic limestone, moderate gray skeletal rock, is wackestone with limestone and siltstone clasts, irregular lens, sponges, very rough, brachiopods (*Meekella*, *Neospirifer*, *Petalichonetes*); conodont sample at 4.5 ft.	2.5	6
1. Limestone and chert, interbedded, silty lime mudstone and rusty siltstone chert, platy and wavy bedded.	3.5	3.5

Section RC VI (Fig. 3)

East shore south flank to Clay Slide.

	Thickness (ft)	Curve letter
Road Canyon Formation		
6. Limestone, silty, sandy, skeletal lime mudstone, thin-bedded with sponges and chert replacements of sponges, chert among bedding planes and joints; conodont sample at 67 ft.	13	43
5. Limestone, silty skeletal lime mudstone with crinoid debris, channeled, brachiopods, platy bedded, barren conodont sample at 52 ft.	20	54
4. Limestone, skeletal lime mudstone with chert as bedding planes, some bedded chert thin-bedded, coarse debris, corals, gastropods, brachiopods (*Derbyia*) at 35 ft; conodont sample at 30.5 ft.	21	54
3. Limestone, silty, skeletal wackestone with chert nodules, in massive irregular lens ranging from 2 to 15 ft thickness within 30 ft of section; conodont sample at 30 ft.	3	33
2. Limestone, silty, skeletal lime mudstone thin and 1 inch sparse fine skeletal wackestone lenses.	13	30
1. Limestone, silty, skeletal lime mudstone with thin rusty chert along bedding planes, thin to medium bedded with thin silty limestone forming wavy mudbeds, crinoid columnals, corals, brachiopods (*Echinaria*, *Rhombella*, others). Forms base of cliff above Clay Slide, barren conodont sample at 9 ft.	19	33

Section RC VII (Fig. 4)

South face on hill facing Hess Canyon, due north of the west end of the hill that lies just north of the Hess house, the east end of the bedrock that contains the type section of the Road Canyon Formation.

	Thickness (ft)	Curve letter
Wood Formation		
11. Limestone, skeletal packstone with corals, sponges, crinoid columnals, brachiopods, and limestone clasts; conodont sample at top, 130 ft.	1	130
10. Siltstone, orange, thin-bedded with few thin carbonate beds.	16	129
Road Canyon Formation		
9. Limestone, light-brownish-gray, skeletal wackestone with carbonate clasts up to boulder size, thick to massive-bedded, sponges, corals, brachiopods (*Derbyia*, *Echinaria*, *Rhombella*, *Meekella*, patches of *Cleiothyridina*); conodont samples at 112, 89 ft.	28	113
8. Mostly covered, apparently orange siltstone.	11	85
7. Limestone, light-tan-gray, pelsal packstone with thin rusty chert stringers, blocky, medium-bedded; corals, fusulinids, siltstone breaks; conodont sample at 70 ft.	14	74
6. Covered, probably orange siltstone.	7	60
5. Limestone, light-tan, sandy, skeletal packstone, massive, large *Cleiothyridina* banks, large crinoid columnals, barren conodont sample at 49 ft.	17	53
4. Limestone, brown-orange, skeletal wackestone, massive, large *Cleiothyridina* bulbs, either one or several piled up; corals, sponges, fusulinids, shaftlimit clasts; crinoid samples at 35, 27 ft; barren samples at 34, 29, 18 ft.	19	36
3. Limestone, tans or light-brownish-gray to orange, silty skeletal packstone, crinoid columnals, sponges, corals, brachiopods (*Composita*, *Rhombella*, *N. richthofeni*); most fossils broken; conodont sample at 11.5 ft.	11.5	17
2. Covered, possibly yellow siltstone.	1.5	5.5
1. Limestone, medium-brownish-gray, skeletal wacke- to packstone, a small mound with sponges, corals, bryozoans, brachiopods (*Composita*, *Cleiothyridina*, *Meekella*, *Neospirifer*, *N. richthofeni*); Spirorbis-like; conodont sample at 1.5 ft.	2	4

Section RC VIII (Fig. 4)

Williams Bar-Crescent Ranch (formerly Appel Ranch) outcrop of the Old Word Ranch site, near USNM localities 703a, c, of Cooper and Grant (1972).

	Thickness (ft)	Curve letter
Word Formation		
7. Limestone, silty lime mudstone, fairly thin-bedded, some cross-laminated, blocky weathering, chert stringers along some bedding planes; few fossils; barren conodont sample at 54 ft.	17	76
6. Covered, thin-bedded yellow siltstone.	18	59
5. Siltstone, yellow, thin-bedded with stringers of silty lime mudstone; barren conodont sample at 39 ft.	6	41
4. Covered, probably yellow siltstone.	4	35
Road Canyon Formation		
3. Limestone, skeletal pelsal wacke- to packstone weathering as rounded carbonate mounds, abundant fossils with clusters of richthofenids (*Cyclacanthia*), *Echinaria*, omegrey, and many other brachiopods; conodont samples at 30.5, 29.5, 27.5, 26 ft; barren sample at 28.5 ft.	6	31
Cathedral Mountain Formation		
2. Mostly covered, yellow siltstone at 20 to 25 ft.	23	25
1. Limestone, light-brownish-gray, crinoidal packstone with fusulinids and few brachiopods, graded with large crinoid columnals at base, more finely broken crinoid debris upward, a wackestone at top; conodont sample at 1.5 ft.	2	2

Section RC IX (Fig. 5)

East flank of Sullivan Peak, on lower slopes directly under the peak itself.

	Thickness (ft)	Curve letter
Wood Formation		
18. Limestone, skeletal wackestone, about 40 percent angular carbonate fragments and 60 percent lime mud (all is Wood siltstone above this bed); conodont sample at 291 ft.	2	292
17. Partly covered, yellow siltstone.	8	290
16. Limestone, skeletal wacke- to packstone, thick-bedded, pelsals, sponges, mudstone pebbles; conodont sample at 281.5 ft.	2	282
15. Partly covered, platy yellow siltstone.	6	280
Road Canyon Formation		
14. Limestone, light-brownish-gray, packstone with carbonate clasts, massive-bedded; corals, sponges, fusulinids, brachiopods (*Cleiothyridina* fragments, *Hustedia*); conodont sample at 272.5 ft.	5	274
13. Partly covered, platy yellow siltstone.	11	267
12. Limestone, medium-brownish-gray, fine-grained pack- to grainstone, medium- to thin-bedded with angular carbonate fragments.	1	256
11. Limestone, medium-brownish-gray, packstone, massive-bedded, angular carbonate clasts of sand to gravel size; chert, corals, cherttified sponges, fusulinids, brachiopods (*Cleiothyridina*); conodont sample at 251.5 ft.	6	255
10. Limestone and siltstone, silty skeletal lime mudstone interbedded with siltstone, black chert nodules; corals, fusulinids, and brachiopods (*Meekella*); conodont sample at 246 ft.	12	247
9. Siltstone, yellow with limestone stringers, lowermost bed is limestone with chert on upper bedding surface; barren conodont sample at 195 ft.	44	235
8. Limestone and siltstone, interbedded, both		

Section RC X (Fig. 4)

West side of Hill 9611, about 1 mi (2 km) SSW of Cornell and Grant (1972).

Road Canyon Formation

9. Siltstone and limestone, moderate yellowish-brown, sandy calcareous siltstone strata (10") alternating with skeletal limestone wackes- to packstone (as much below) and yellowish-brown, silty lime mudstone	8	80
8. Limestone, pale yellowish-brown, skeletal packstone with abundant crinoid columnals; conodont sample at 52 ft	11	55
7. Siltstone, yellowish-orange, thin-bedded	9	42
6. Limestone, pale yellow-orange, skeletal wacke- to packstone with abundant crinoid columnals, scattered chert nodules	4	33
5. Limestone, pale yellowish-brown, skeletal packstone with abundant brachiopods and crinoid columnals; conodont sample at 29 ft	3	29
4. Limestone, pale yellowish-brown, slightly recrystallized, skeletal wacke- to packstone with abundant crinoid columnals, scattered chert nodules	13	26
3. Covered	2	13
2. Limestone, recrystallized, skeletal wacke- to packstone with abundant crinoid columnals, beds mottling and swelling	8	11
1. Limestone, pinkish-gray, recrystallized, skeletal wackestone, recrystallization due to close proximity to Hess Hunt Boundary fault; conodont sample at 1 ft	3	3

Section RC XI (Fig. 4)

South face of knoll that is USSM vicinity 7(2), Cornell and Grant, 1972; includes north-facing highway

Road Canyon Formation

2. Limestone, heavily exposed on cliffside, light brownish-gray, skeletal wackestone	15	15
1. Limestone, light brownish-gray, skeletal wackestone, massive-bedded with rare, low-relief stromatolite bioherms throughout but more common near top; lime mudstone clasts in lower 2.5 ft; conodont samples at 8.5, 13.5, 15.5 ft	15	15

Section RC XII (Fig. 4)

Limestone ledge east of Apex Tank, about 1 USSM location 7(2m), 7(2nw), 7(2q), 7(2ne), 7(2se), 7(2sw), and 7(3q), earlier to Fig. 22 in Cooper and Grant (1969, 1972)

Road Canyon Formation

1. Limestone, pale-yellowish-brown, skeletal wackestone, medium- to thick-bedded; conodont samples at 5.5, 11.5 ft	30	30

REFERENCES

Baird, M.R., 1975, Conodont biostratigraphy of the Kaibab Formation, eastern Nevada and western Utah: unpublished Master's thesis, Ohio State University, 71 p.

Behnken, F.H., 1975, Leonardian and Guadalupian (Permian) conodont biostratigraphy in western and southwestern United States: Journal of Paleontology, v. 49, p. 284–315.

Behnken, F.H., Wardlaw, B.R., and Stout, L.N., 1986, Conodont biostratigraphy of the Permian Meade Peak Phosphatic Shale Member, Phosphoria Formation, southeastern Idaho: Contributions to Geology, University of Wyoming, v. 24, no. 2, p. 169–196.

Bender, Hans, and Stoppel, Dieter, 1965, Perm-Conodonten: Geologisches Jahrbuch, v. 82, p. 331–364.

Clark, D.L., and Behnken, F.H., 1971, Conodonts and biostratigraphy of the Permian, in Sweet, W.C., and Bergström, S.M., eds., Symposium on biostratigraphy: Geological Society of America Memoir 127, p. 415–439.

——1979, Evolution and taxonomy of the North American Upper Permian Anchignathodus–Sweetognathus complex: Journal of Paleontology, v. 53, p. 263–275.

Clark, D.L., and Ethington, R.L., 1962, Survey of Permian conodonts in western North America: Brigham Young University Geology Studies, v. 9, pt. 2, p. 107–114.

Clark, D.L., and Mosher, L.C., 1966, Stratigraphic, geographic and evolutionary development of the conodont genus Sweetognathus: Journal of Paleontology, v. 40, p. 1376–1394.

Cooper, G.A., and Grant, R.E., 1964, New Permian stratigraphic units in Glass Mountains, West Texas: American Association of Petroleum Geologists Bulletin, v. 48, p. 2342–2355.

——1966, Permian brachiopods of Glass Mountains, West Texas: U.S. Geological Survey Bulletin 1244-E, 63 p.

——1972, Permian brachiopods of West Texas, 1: Smithsonian Contributions to Paleobiology, no. 14, 230 p.

——1975, Dating and correlating the Permian of the Glass Mountains in Texas, in Logan, Alan, and Hills, L.V., eds., The Permian and Triassic systems and their mutual boundary: Canadian Society of Petroleum Geologists Memoir 2, p. 522–549.

Ito, Hisaharu, 1981, Permian conodont biostratigraphy of Japan: Palaeontological Society of Japan Special Paper 24, 90 p.

King, P.B., 1931, The geology of the Glass Mountains, Texas, pt. 1, Descriptive geology: University of Texas Bulletin, no. 3038, 167 p.

——1942, Permian of West Texas and southeastern New Mexico: American Association of Petroleum Geologists Bulletin, v. 26, p. 535–763.

Kozur, Heinz, 1975, Beitrage zur Conodontenfauna des Perm (Contributions to the conodont fauna of the Permian): Geologisch-Palaontologische Mitteilungen Innsbruck, v. 5, no. 4, p. 1–44.

——1978, Beitrage zur Stratigraphie des Perm, Teil II: Die Conodonten-Chronologie des Perms (Contributions to the stratigraphy of the Permian, Part II: The conodont chronology of the Permian): Freiberger Forschungsheft, v. C 334, p. 45–161.

Kozur, Heinz, and Mostler, Helfried, 1979, Anisaxonitella n. gen. und Merrillina n. gen., zwei neue Conodontengattungen aus dem Perm (Anisaxonitella n. gen. and Merrillina n. gen., two new conodont genera from the Permian): Geologisch-Palaontologische Mitteilungen Innsbruck, v. 8, no. 7, p. 259–282.

Kozur, Heinz, Mostler, Helfried, and Rahimi-Yazd, Ali, 1975, Beitrage zur conodontenfauna permotriadischer schichtfolgen Teil II: Neue conodonten aus dem Oberen Perm und der Trias des Iran und Nordost-Afghanistan (Contributions to the conodont fauna of Permo-Triassic sequences, Part II: New conodonts from the Upper Permian and the basal Triassic of North and Central Iran): Geologische Palaontologische Mitteilungen Innsbruck, v. 5, no. 3, p. 1–23.

Mamay, S.H., Miller, J.M., and Rohr, D.M., 1984, Late Leonardian plants from West Texas: The youngest plant megafossils in North America: Science, v. 223, p. 279–281.

Merrill, G.K., and Powell, R.L., 1980, Paleobiology of juvenile cuspinate conodonts from the Drum Limestone (Pennsylvanian, Missourian), Kansas City areas and its bearing on apparatus ontogeny: Journal of Paleontology, v. 54, p. 1058–1074.

Miller, A.K., and Furnish, W.M., 1957, Ammonoids of the basal Word Formation, Glass Mountains, West Texas: Journal of Paleontology, v. 31, p. 1052–1056.

Rohr, D.M., Miller, J.M., Davis, R.A., and Mamay, S.H., 1982, An Upper Permian (Guadalupian) flora from the Glass Mountains area, Brewster County, West Texas: Geological Society of America Abstracts with Programs, v. 14, no. 7, p. 662.

Rohr, D.M., Davis, R.A., Mamay, S.H., and Miller, J.M., 1987, Leonardian plant-bearing beds from the Del Norte Mountains, West Texas, in Cromwell, Dave, and Mazzullo, L.J., eds., The Leonardian facies in W. Texas and S.E. New Mexico and guidebook to the Glass Mountains, West Texas: 1987 Permian Basin Section, Society of Economic Paleontologists and Mineralogists Publication 87-27, p. 67–68.

Sweet, W.C., 1977a, Hindeodus, in Ziegler, Willi, ed., Catalogue of conodonts, E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, v. 3, p. 203–224.

——1977b, Xaniognathus Bischoff, in Ziegler, Willi, ed., Catalogue of conodonts, E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, v. 3, p. 291–295.

Wang, Zhi-hao, 1978, Permian-Lower Triassic conodonts of the Langshan area, southern Shaanxi: Acta Palaeontologica Sinica, v. 17, p. 213–227.

Wardlaw, B.R., 1980, Middle-Late Permian paleogeography of Idaho, Montana, Nevada, Utah, and Wyoming, in Fouch, T.D., and Magathan, E.R., eds., Paleogeography of west-central United States: West-central United States Paleogeography Symposium 1, Society of Economic Paleontologists and Mineralogists, Rocky Mountain Section, p. 353–361.

Wardlaw, B.R., and Collinson, J.W., 1978, Stratigraphic relations of the Park City Group (Permian) in eastern Nevada and western Utah: American Association of Petroleum Geologists Bulletin, v. 62, p. 1371–1384.

——1979, Youngest Permian conodont faunas from the Great Basin and Rocky Mountain regions, in Sandberg, C.A., and Clark, D.L., eds., Conodont biostratigraphy of the Great Basin and Rocky Mountains: Brigham Young University Studies, v. 26, pt. 3, p. 251–263.

——1984, Conodont paleoecology of the Permian Phosphoria Formation and related rocks of Wyoming and adjacent areas: Geological Society of America Special Paper 196, p. 263–284.

——1986, Paleontology and deposition of the Phosphoria Formation: Contributions to Geology, University of Wyoming, v. 24, no. 2, p. 107–142.

Wardlaw, B.R., and Grant, R.E., 1987, Conodont biostratigraphy of the Cathedral Mountain and Road Canyon Formations, Glass Mountains, West Texas, in Cromwell, Dave, and

he Leonardian facies in W. Texas and
guidebook to the Glass Mountains
rmian Basin Section: Society of Eco-
and Mineralogists Publication 67-27.

d-defined Permian stages, in Permian
s, and stratigraphy: West Texas Geo-
rmian Basin Section of the Society of
gists and Mineralogists Publication

Wilde, G.L., and Todd, R.G., 1968, Guadalupian biostratigraphic relationships and sedimentation in the Apache Mountains region, West Texas, pt. 1, Surface correlations: Symposium and Guidebook, 1968 Field Trip, Permian Basin Section, Society of Economic Paleontologists and Mineralogists, Publication 68-11, p. 10–31.

Youngquist, W.L., Hawley, R.W., and Miller, A.K., 1957, Phosphoria conodonts from southeastern Idaho: Journal of Paleontology, v. 25, p. 356–364.

PLATES 1–4

Contact photographs of the plates in this report are available, at cost, from the U.S. Geological Survey Photographic Library, Federal Center, Denver, CO 80225.

PLATE 1

NEOGONDOLELLA

(All figures ×60)

Figures 1–7. *Neogondolella serrata* (Clark and Ethington)
1. Oblique upper view. Pa element. USNM 406891, W81169.
2. Upper view. Pa element. USNM 406892. W81169.
3. Upper view. Pa element. USNM 406893. W81169.
4. Oblique upper view. Pa element. USNM 406894. W81176.
5. Upper view. Pa element. USNM 406895. W81176.
6. Oblique upper view. Pa element. USNM 406896. W81176.
7. Upper view. Pa element. USNM 406897. W81174.

10–13. *Neogondolella serrata* (Clark and Ethington)
10. Oblique upper view. Pa element. USNM 406898. W81176.
11. Upper view. Pa element. USNM 406899. W81176.
12. Oblique upper view. Pa element. USNM 406900. W81176.
13. Oblique upper view. Pa element. USNM 406901. W81169.

17–18. *Neogondolella serrata* (Clark and Ethington)
17. Oblique upper view. Pa element. USNM 406902. W81176.
18. Upper view. Pa element. USNM 406903. W81176.

8–9. *Neogondolella idahoensis* (Youngquist, Hawley, and Miller)
8. Oblique upper view. Pa element. USNM 406904. W81161.
9. Oblique upper view. Pa element. USNM 406905. W81161.

14–16. *Neogondolella idahoensis* (Youngquist, Hawley, and Miller)
14. Upper view. Pa element. USNM 406906. W81179.
15. Upper view. Pa element. USNM 406907. W81161.
16. Oblique upper view. Pa element. USNM 406908. W81179.

19–22. *Neogondolella idahoensis* (Youngquist, Hawley, and Miller)
19. Upper view. Pa element. USNM 406909. W81161.
20. Oblique upper view. Pa element. USNM 406910. W81161.
21. Oblique upper view. Pa element. USNM 406911. W81161.
22. Oblique upper view. Pa element. USNM 406912. W81179.



NEOGONDOLELLA

PLATE 2

HINDEODUS AND *NEOGONDOLELLA*

(All figures ×60)

Figures 1–15. *Hindeodus excavatus* (Behnken).
1. Inner view, Sc element, USNM 406913, W81176.
2. Inner view, Sc element, USNM 406914, W81176.
3. Inner view, Pb element, USNM 406915, W81176.
4. Inner view, Pb element, USNM 406916, W81176.
5. Lateral view, Pa element, USNM 406917, W81176.
6. Lateral view, Pa element, USNM 406918, W81176.
7. Inner view, Sc element, USNM 406919, W81176.
8. Posterior view, M element, USNM 406920, W81176.
9. Lateral view, M element, USNM 406921, W81176.
10. Posterior view, Sb element, USNM 406922, W81176.
11. Posterior view, Sb element, USNM 406923, W81176.
12. Outer view, Sc element, USNM 406924, W81176.
13. Inner view, Sc element, USNM 406925, W81176.
14. Posterior view, Sb element, USNM 406926, W81176.
15. Outer view, Sc element, USNM 406927, W81176.

16–19. *Neogondolella serrata* (Clark and Ethington).
16. Lower view, Pa element, USNM 406928, W81174.
17. Lower view, Pa element, USNM 406929, W81176.
18. Lower view, Pa element, USNM 406930, W81169.
19. Lower view, Pa element, USNM 406931, W81169.

23–25. *Neogondolella serrata* (Clark and Ethington).
23. Lower view, Pa element, USNM 406932, W81176.
24. Lower view, Pa element, USNM 406933, W81176.
25. Lower view, Pa element, USNM 406934, W81176.

20–22. *Neogondolella idahoensis* (Youngquist, Hawley, and Miller).
20. Lower view, Pa element, USNM 406935, W81179.
21. Lower view, Pa element, USNM 406936, W81161.
22. Lower view, Pa element, USNM 406937, W81161.

26–28. *Neogondolella idahoensis* (Youngquist, Hawley, and Miller).
26. Lower view, Pa element, USNM 406938, W81161.
27. Lower view, Pa element, USNM 406939, W81179.
28. Lower view, Pa element, USNM 406940, W81161.



PLATE 3

NEOSTREPTOGNATHODUS, HINDEODUS, SWEETOGNATHUS, AND SWEETINA

[All figures ×60, except where noted otherwise]

Figure 1. *Neostreptognathodus newelli* Wardlaw and Collinson.
1. Upper view, Pa element, USNM 406941, W81228.

2. *Neostreptognathodus clinei* Behnken.
2. Upper view, Pa element, USNM 406942, W81243.

3–11. *Hindeodus excavatus* (Behnken).
3. Upper view, Pa element, USNM 406943, W81176.
4. Oblique upper view, Pa element, USNM 406944, W81176.
5. Lateral view, Pa element, USNM 406945, W81176.
6. Lateral view, Pa element, USNM 406946, W81176.
7. Lateral view, Pa element, USNM 406947, W81176.
8. Lateral view, Pa element, USNM 406948, W81176.
9. Lateral view, Pa element, USNM 406949, W81176.
10. Lateral view, Pa element, USNM 406950, W81176.
11. Lateral view, Pa element, USNM 406951, W81176.

12–17. *Sweetognathus iranicus* Kozur, Mostler, and Rahimi-Yazd.
12, 16. Upper view, Pa element, USNM 406952, W81176, ×60 and enlarged, ×300, showing surface pustules.
13, 17. Upper view, Pa element, USNM 406953, W81176, ×60 and enlarged, ×300, showing surface pustules.
14. Upper view, Pa element, USNM 406954, W81194.
15. Upper view, Pa element, USNM 406955, W81200.

18–25. *Sweetina hottica* (Bender and Stoppel).
18. Lateral view, Sa element, USNM 406956, W81165.
19. Posterior view, M element, USNM 406957, W81176.
20. Posterolateral view, M element, USNM 406958, W81176.
21. Posterolateral view, Sb element, USNM 406959, W81162.
22. Inner view, Sc element, USNM 406960, W81176.
23. Inner view, Sc element, USNM 406961, W81176.
24. Lateral view, Sc element, USNM 406962, W81176.
25. Posterior view, M element, USNM 406963, W81165.



PLATE 4

XANIOGNATHUS, NEOSTREPTOGNATHODUS, AND NEOGONDOLELLA

[All figures ×60]

Figures 1–22. *Xaniognathus abruptus* (Clark and Ethington):
1. Posterior view, Sa element, USNM 406964, W81239.
2. Oblique lateral view, Sa element, USNM 406965, W81176.
3. Oblique lateral view, Sa element, USNM 406966, W81176.
4. Posterior view, M element, USNM 406967, W81176.
5. Anterior view, M element, USNM 406968, W81176.
6. Posterolateral view, M element, USNM 406969, W81176.
7. Posterior view, M element, USNM 406970, W81176.
8. Posterior view, Sb element, USNM 406971, W81176.
9. Posterior view, Sb element, USNM 406972, W81176.
10. Posterolateral view, Sb element, USNM 406973, W81176.
11. Anterolateral view, Sb element, USNM 406974, W81176.
12. Lateral view, Pa element, USNM 406975, W81176.
13. Lateral view, Pa element, USNM 406976, W81176.
14. Lateral view, Sc element, USNM 406977, W81239.
15. Lateral view, Sc element, USNM 406978, W81176.
16. Lateral view, Sc element, USNM 406979, W81176.
17. Lateral view, Pa element, USNM 406980, W81176.
18. Lateral view, Pa element, USNM 406981, W81176.
19. Inner view, Pb element, USNM 406982, W81176.
20. Inner view, Pb element, USNM 406983, W81176.
21. Inner view, Pb element, USNM 406984, W81176.
22. Inner view, Pb element, USNM 406985, W81176.

23. *Neostreptognathodus* sp.
23. Inner view, Pb element, USNM 406986, W81176.

24–25. *Neogondolella idahoensis* (Youngquist, Hawley, and Miller):
24. Upper view, Pa element, USNM 406987, W81181.
25. Upper view, Pa element, USNM 406988, W81181.



Chapter B

The Rugose Coral *Ankhelasma*—Index to Viséan (Lower Carboniferous) Shelf Margin in the Western Interior of North America

By WILLIAM J. SANDO, E. WAYNE BAMBER, and BARRY C. RICHARDS

Predictions of Carboniferous shelf-margin locations using the living zone of a rugose coral

U.S. GEOLOGICAL SURVEY BULLETIN 1895
SHORTER CONTRIBUTIONS TO PALEONTOLOGY AND STRATIGRAPHY

CONTENTS

Abstract B1

Introduction B1

Acknowledgments B2

Paleoecologic Setting B3

Western United States B3

Western Canada B3

Stratigraphic, Lithographic, and Ecologic Distribution of Ammonites B4

Western United States B4

Western Canada B11

Interval Shell Margin Extinctions B14

Western United States B14

Western Canada B17

Conclusions B17

References Cited B17

Appendixes A–D

A. Ammonite Localities in the United States B19

B. Coastal Points for Chronostratic Lithofacies Profiles across Known Upper Shell Margin in the United States B24

C. Ammonite Localities in East-Central British Columbia, Canada B24

D. Coastal Points for Chronostratic Lithofacies Profiles across Inferred Upper Shell Margin in Manitoba B29

FIGURES

1. Map of western North America showing area of Lower Cretaceous sedimentary rocks in Western Interior region and Lower Cretaceous paleotectonic elements B1
2. Photographs of specimens of Ammonites Sands, 1867, from the Western Interior region of North America B2
3. Map showing Cretaceous igneous and metamorphic elements, Western Canada Shield B4
4. Cartographic image of Western Canada in Western Interior region of the United States showing locations of Ammonites occurrences B5
5. Temporal and stratigraphic distribution of Ammonites in the Western United States B6
6. Map showing early Vostok (Campanian Zones 12 and 13) paleogeography of a part of the Western Interior basin in the United States and Ammonites occurrences in relation to shell margin and Disphotic B8
7. Map showing early middle Vostok (upper Campanian Zone 14) paleogeography of a part of the Western Interior basin in the United States and Ammonites occurrences in relation to shell margin and Disphotic B9

8. Chronometric, lithofacies profiles across Visean shelf margin in the Western United States B10
9. Map showing present nonpalinspastic facies distribution of Visean (Composite Zones 12 to 14) deposits, east-central British Columbia and Ankhelasma occurrences in relation to shelf margin B12
10. Stratigraphic cross section showing Ankhelasma occurrences and depositional environments in Rundle Group and Besa River Formation, study area, east-central British Columbia B13
11. Diagram showing generalized depositional model of Canadian Early Carboniferous carbonate platform B14
12. Temporal and stratigraphic distribution of Ankhelasma in east-central British Columbia B15
13. Chronometric, lithofacies profiles from Ankhelasma localities in southwestern Canada to sections in northwestern Montana to inferred shelf margin B16

TABLE

1. Nonpalinspastic distances of Ankhelasma occurrences landward from known Visean shelf margin in Western Interior region of the United States B7

The Rugose Coral *Ankhelasma*—Index to Viséan (Lower Carboniferous) Shelf Margin in the Western Interior of North America

By William J. Sando,[1] E. Wayne Bamber,[2] *and* Barry C. Richards[3]

Abstract

Ankhelasma, a distinctive, endemic, solitary rugose coral, lived in a belt presently no more than 75 kilometers wide, almost exclusively on the landward side of the Visean shelf margin in the Western Interior region of western North America. A model of the distribution of this coral, based on known nonpalinspastic distances to the shelf margin in tracks of early Visean and early middle Visean age, suggests that the living zone of this coral was probably no more than approximately 30 kilometers wide. The genus originated either in northeastern Utah or southwestern Canada at the beginning of the early Visean (Composite Zone 12) and migrated both northward and southward along the shelf margin, dying out by the end of early, late Visean (Composite Zone 14) time, a phylogenetic life span of about 8 million years. Occurrences of this coral in northwestern Montana and east-central British Columbia adjacent to areas where the Visean depositional record has been removed by subsequent erosion are helpful in determining or confirming inferred locations for the shelf margin in these areas.

Figure 1. Western North America showing area of Lower Carboniferous sedimentary rocks in Western Interior region stippled; and Lower Carboniferous paleotectonic elements. Compiled from Sproule (1950, fig. 1; Poole and Sandberg (1977, fig. 1; Sando (1985, fig. 1; and Richards and others (in press). Triangles indicate areas that have one or more occurrences of *Ankhelasma*.

INTRODUCTION

During the Early Carboniferous, the Western Interior region of North America was occupied by an epeiric sea that extended from arctic Canada to northern Mexico (fig. 1). *Ankhelasma* Sando, 1961 (fig. 2), is a small, curved-conical, solitary rugose coral, endemic to the Western Interior province of Sando and others (1975, 1977), that is distinguished by the extraordinary development of five septal plates in the mature (ephebic) stage. The presence of these plates facilitates recognition in transverse section, even without laboratory preparation. The genus is represented by one nominal species, *A. typicum* Sando, 1961, in the United States and Canada and by at least one unnamed species in Canada. *Ankhelasma* probably evolved from the widespread

Manuscript approved for publication June 27, 1985.

[1] U.S. Geological Survey, U.S. National Museum of Natural History, Washington, DC 20560.
[2] Geological Survey of Canada, 3303-33rd St. NW., Calgary, Alberta, Canada T2L 2A7.

genus *Lithostrotion* (*Homalophyllites* of earlier reports; see Sando, 1961, p. 67), a coral that occurs commonly in the sequence below the lowest occurrence of *Ankhelasma* and is sparse in the lowermost part of the range of *Ankhelasma* (Sando and Bamber, 1985).

Preliminary studies of the stratigraphic and geographic distribution of *Ankhelasma* suggested that it was



Figure 2. Specimens of Anthelasma Sandy, 1981 from the Western Interior region of North America. A, B, Calicular and alar views, respectively, ×1, of silicified coralliom of A. typicum, holotype, USNM 120201 from USGS loc. 17722-PC, Brazer Dolomite (member 3), Composite Zone 12, Rex Peak, Crawford Mountains, Utah. C, D, Calicular and alar views, respectively, ×1, of silicified coralliom of A. typicum, hypotype, USNM 424262 from USGS loc. 27068-PC, Monte Cristo Limestone, Yellowpine Member, Composite Zone 34, Potosi Mine, Goodsprings District, Nevada. E, F, Calicular and alar views, respectively, ×1, of silicified coralliom of A. cf. sp., hypotype, CSC 9864 from GSC loc. C-110691, Prophet Formation (member 8), Composite Zone 33 or 34, South Mountain Creek, Pine Pass map area, east-central British Columbia. G, Transverse thin section, ×10, of A. typicum, hypotype, USNM 424263 from USGS loc. 27074-PC, Brazer Dolomite (member 2), Composite Zone 12, Trail Hollow Canyon Mountains, Utah. H, Thin section of Anthelasma bearing fossil rudstone, A. typicum and associated fossils in paleocurrent (×2, same locality as C).

restricted to a narrow geographic zone related to the edge of a broad carbonate shelf (shelf margin) during the early to early middle Visean in the Western Interior region of the United States. A similar distribution pattern was noted in western Canada, where exposures of the shelf margin are extensive but less completely sampled.

Because shelf margins define depositional strike along basin flanks and influence the character and distribution of petroleum-reservoir rocks and seals, they are of prime economic importance (Rose, 1976, p. 449–458). Accurate delineation of Carboniferous shelf margins is important to the preparation of regional paleogeographic maps, which can be used to analyze depositional history. Mapping of shelf margins also can be useful for measuring later structural offsets and translocations. This paper documents a geographic distribution model for Anthelasma and uses this model to predict or confirm shelf-margin locations in areas where the depositional record has been removed or obscured by later events.

Acknowledgments

Early field work by the senior author on the stratigraphy of Anthelasma was done largely with I.T. Dutro, Jr., whose companionship and contributions to the stratigraphy are gratefully acknowledged. Ten years of joint field studies with C.A. Sandberg and R.C. Gutschick laid the foundation for sophisticated analyses of the shelf margin; the creative and inspiring conceptual contributions of these colleagues strongly influenced many of the ideas expressed in this paper. F.E. Bierwagen, J.A. Dockal, M.R. Mudge, C.A. Sandberg, and Betty Skipp contributed samples of Anthelasma to the senior author. Shell Canada Limited supplied specimens and information from several localities in Canada. Consultant identifications and zonal determinations were made by C.A. Sandberg for samples from the United States, and A.C. Higgins was responsible for the same information on Canadian samples. Identifications of foraminifers and algae and zonal determinations on samples from the United States and Canada were made by B.L. Mamet.

PALEOTECTONIC SETTING

Western United States

Previous studies of Devonian and Carboniferous geology in the Western United States (Poole, 1974; Nilsen, 1977; Poole and Sandberg, 1977) showed that the Western Interior basin was located at the edge of the North American protocontinent, and that the basin was affected by Late Devonian and Early Carboniferous plate collisions (Antler orogeny). A broad, shallow, epicontinental sea, located on the craton (cratonic platform), was bounded on the east by the Transcontinental arch, a positive area of Precambrian crystalline rocks and lower Paleozoic sedimentary rocks that formed a peninsula during the Early Carboniferous and separated the Western Interior and Eastern Interior basins (Fig. 1) (Gutschick and Sandberg, 1983). An island chain composed of uplifted early Paleozoic rocks was formed in the Antler orogenic belt during plate collision and bounded the Western Interior basin on the west. Plate collision also produced a deep trough (Antler foreland basin) between the Antler orogenic belt and the stable cratonic platform during the Visean.

Deep-water, terrigenous and carbonate sediments deposited in the Antler foreland basin were separated from shallow-water carbonate and evaporite sediments of the cratonic platform by a narrow break in depositional slope, called shelf margin by previous workers (Rose, 1976; Sando, 1976; Gutschick and others, 1980; Sandberg and others, 1982; Gutschick and Sandberg, 1983). The paleographic product of the Mississippian shelf margin changed through time, but the shelf margin was always close to the craton margin. Anthelasma has been found only in a narrow belt that follows the tectonic zone of Lateralus thinning, approximately coincident with the craton margin and Visean shelf margin, from southeastern Nevada to northwestern Montana (Fig. 1).

Western Canada

Canadian occurrences of Anthelasma are of the same age as those in the United States and occupy similar geographic positions with respect to the craton margin and Visean shelf margin (Figs. 1 and 3). Strata containing Anthelasma were deposited in the southwestern part of the Peace River embayment and in the Prophet trough to the west and south. Although the distribution pattern is similar, there are still some uncertainties concerning exact relationships with paleotectonic elements recognized in the United States.

The name Prophet trough was introduced by Richards and others (in press) for an extensive, northwest-trending, downwarped and downfaulted belt that developed on the western margin of the North American protocontinent during the latest Devonian and Carboniferous. The trough was continuous with the Antler foreland basin and extended from southeastern British Columbia into the northern part of the Yukon Territory. A broad hinge zone, marking a point at which water depths and sedimentation rates increased rapidly basinward, formed the boundary between the Prophet trough and the cratonic platform to the east. The western boundary of the trough was an elevated rim, at least locally subaerially exposed during the Early Carboniferous. The Antler orogenic belt of the United States, or a related orogenic belt, may have extended at least as far north as the southern part of the Peace River embayment, thereby forming part of the western rim of the southern Prophet trough (Richards and others, in press). The origin of the trough is not well established. Some workers (Templeman-Kluit, 1979; Gordey and others, 1987; Struik, 1987) believe that the trough is either partly or predominantly an extensional feature resulting from Late Devonian and Early Carboniferous rifting of western North America. Extension, recorded by extensive block faulting, did occur in the central part of the Prophet trough and in the Peace River embayment (Richards and others, in press). However, tectonic and sedimentologic data summarized by Richards and others (in press) suggest that the Antler orogeny or a related compressional event influenced the marked Tournaisian and Visean subsidence that occurred in the southern part of the trough.

In southwestern Alberta and northeastern British Columbia, the Peace River embayment, a broad reentrant into the cratonic platform, opened westward into the Prophet trough (Figs. 1 and 3). The depositional and structural axis of this arcuate embayment had an easterly trend and coincided approximately with the axis of the Late Devonian Peace River arch. Extensive block faulting, accompanied by regional flexural subsidence, produced the embayment, which is characterized by an anomalously thick Lower Carboniferous sequence and by numerous northeasterly and northwesterly striking synsedimentary and postdepositional normal faults. Evidence for westward shallowing during the Tournaisian and Visean indicates that a slightly positive area, possibly a series of islands or a peninsula, was at least periodically present in the southwestern part of the Peace River embayment where strata containing Anthelasma were deposited. Part of this northwest-trending positive area, which extended southeastward of the embayment, were emergent, eroded, and exposed to subaerial erosion during the earliest middle Tournaisian (earliest Tn2) to Serpukhovian. All

Figure 1. Carboniferous deposits and paleotectonic elements, Western Canada basin (after Richards and others, in press). Study area contains all known Canadian occurrences of *Ankhelasma*; see fig. 4.

known Canadian occurrences of *Ankhelasma* are in carbonate rocks deposited on or near the positive region.

STRATIGRAPHIC, GEOGRAPHIC, AND ECOLOGIC DISTRIBUTION OF *ANKHELASMA*

Western United States

In the United States, *Ankhelasma* occurs at 17 surface localities in a narrow belt (Ankhelasma belt) along the western margin of the cratonic platform (fig. 4) (see appendix A for complete descriptions of localities). The spacing of occurrences along the strike of the Ankhelasma belt seems to be determined mainly by availability of outcrops of rocks potentially bearing *Ankhelasma*; this relation is most notable in the long stretch of land having no occurrences from the southwestern corner of Utah to northeastern Utah, where rocks potentially bearing *Ankhelasma* do not crop out. Consequently, *Ankhelasma* is not recorded from this area, although no subsurface cores have been examined by the writers.

More than 80 years of intensive collecting of corals by the senior author outside the area herein designated Ankhelasma belt, as well as hundreds of collections sent to the senior author for study by other geologists, have failed to reveal even a single specimen of the genus. Moreover, examination of hundreds of collections of Lower Carboniferous rocks made by geologists of the U.S. Geological Survey during the past 80 years revealed only two occurrences of this unique coral, and these were both in the narrow belt specified by more recent field work.

Ankhelasma occurs in two stratigraphic intervals in the United States (fig. 4). In northeastern Utah, western Wyoming, southeastern Idaho, and western Montana, the occurrences are all dated as early Visean (Composite Zones 12 and 13 of Sando, 1985) by foraminifers, corals, and conodonts (localities 2–8, 10–13). In south-central Idaho, a poorly dated debris-flow occurrence (locality 9) is also thought to be from the same interval. In southeastern Nevada and northwestern Arizona (localities 1–3), the occurrences are of early middle Visean age (Composite Zone 14 of Sando, 1985).

A more detailed analysis of stratigraphic distribution (fig. 5) shows that the earliest known occurrences of *Ankhelasma* in the United States are at the base of Composite Zone 12 at localities 4 and 5 in the Uinta Mountains of northeastern Utah. The genus ranges through Composite Zone 12 well into Composite Zone 13 along the Ankhelasma belt to the northeast in western Wyoming, southeastern Idaho, and western Montana (localities 6–13).


ANTLER OROGENIC BELT
FORELAND BASIN
CRATONIC PLATFORM
TRANSCONTINENTAL ARCH
EXPLANATION
VISEAN DEPOSITIONAL AREAS
Outcrop of older rocks where Visean rocks were removed by subsequent erosion
Visean rocks preserved in surface and subsurface (includes small areas removed by subsequent erosion)
Visean rocks mostly obscured by Cretaceous-Tertiary sedimentary, extrusive, and intrusive rocks
VISEAN POSITIVE AREAS
Visean rocks never deposited (includes some areas removed by subsequent erosion). Dashed lines indicate inferred extensions of positive areas
Post-Visean lateral fault, dashed where inferred. Arrows indicate relative direction of movement
Post-Visean thrust fault, dashed where inferred. Sawteeth on upper plate
Southern limit of Pre-Jurassic truncation
Ankhelasma occurrences
Composite zone 14
Composite zone 12 or 13
Debris flow, Composite zone 12
Area of map
0 100 MILES
0 100 KILOMETERS

Figure 4. Geologic map of Visean rocks in Western Interior region of the United States showing locations of *Ankhelasma* occurrences. Modified from Craig and Connor (1979, pl. 2) and Sandberg and others (1982, fig. 2).

TIME SCALES
1
2
3
4
5
6
7
8
9
10
11
12
13
COMPOSITE BIOZONES
ASSEMBLAGES
DISCORBIDS
CASSIDULINIDS
ELPHIDIIDS
ROSALINIDS
MILIOLIDS
SPIROPLECTAMMINIDS
TEXTULARIIDS
CASSIDULINIDS
ELPHIDIIDS
ROSALINIDS
MILIOLIDS
SPIROPLECTAMMINIDS
TEXTULARIIDS
DISCORB

Shelf Margin 1
MADISON SHELF
BASIN
PEORIA TROUGH
HUBBARBINE RIDE
HIGH
CENTRAL LANDMASS
BASIN
Shelf Margin 1
LEADVILLE REDWALL SHELF
EXPLANATION
Visean depositional record lost or obscured by later events
Visean rocks never deposited or removed by later erosion
Post-basinal facies on highs or rises, later eroded
Open-marine shelf facies
Restricted shelf
Open-marine basin facies
Starved basin facies
Karst facies
Post-Visean lateral fault, dashed where inferred. Arrows indicate relative direction of movement
Post-Visean thrust fault, dashed where inferred. Sawteeth on upper plate
Shelf margin 1. Dashed where approximate, dotted where inferred
Chronostratic cross section profile shown on figs. 9 and 13
Control points in vicinity of shelf margin
Actual
Inferred
Anadelasma occurrences
on Mission River
on Jessica River
0 50 100 MILES
0 50 100 KILOMETERS

Figure 6. Early Visean (Composite Zones 12 and 13) paleogeographs of a part of the Western Interior basin in the United States and Anadelasma occurrences in relation to shelf margin and lithofacies. Modified from Sandberg and others (1982, fig. 9).

Figure 7. Early middle Visean (upper Composite Zone 14) paleogeographs of a part of the Western Interior basin in the United States and Anadelasma occurrences in relation to shelf margin and lithofacies. Modified from Sandberg and others (1982, fig. 11).



Figure 8. Chronostratic lithofacies profiles across Upper shelf margin in the Western United States. Geographic locations of profiles shown on figures 6 and 7. Control points for these profiles are noted and described in appendix B. Ma = million years before present; radiometric scale; CZ = composite zone.

since the base of Composite Zone 12 in two localities (see subsequent discussion in Western Canada section). Evidently, the genus originated either in northwestern Utah and migrated rapidly northward, or it originated in British Columbia or Alberta and migrated rapidly southward during earliest Visean time (Hogarth time unit number 1 or 2; fig. 5).

The lack of outcrops along the shelf margin between northwestern Utah and the southwestern corner of Utah prevents determination of the southern limit of *Anthracoceras* migration during the late early Visean (Composite Zone 13). The genus has not been found in the Redwall Dolomite Member of the Monte Cristo Limestone in southeastern Nevada and southwestern Utah, or in the Mooney Falls Member of the Redwall Limestone in northwestern Arizona, a distribution that suggests that *Anthracoceras* had not reached the southern part of the shelf at this time. However, the occurrence of *Anthracoceras* in the Yellowjacket Member of the Monte Cristo Limestone (upper half of Composite Zone 14) suggests that the genus was by then in the intervening area of no outcrop during the late early Visean (Composite Zone 13).

The early middle Visean (Composite Zone 14) was characterized by uplift and subaerial erosion of the northern Madison shelf area to form the Madison karst plain (Sando, 1988; fig. 7). A sabkha, represented by the McKenzie Canyon Limestone in southwestern Montana and the upper most part of the Monroe Canyon Limestone in southeastern Idaho and western Wyoming, occupied the shore of the karst plain. A tremendous volume of quartz sand derived from the uplifted Transcontinental arch to the east was brought by rivers westward across the karst plain in northwestern Colorado to form the Humbug Formation in northeastern Utah and equivalent rocks in southwestern Wyoming and southeastern Idaho (Sandberg and others, 1983, p. 774). Meanwhile, open marine shelf carbonate deposition continued on the Leadville Redwall shelf in southern Utah, northern Arizona, and southeastern Nevada. By this time, carbonate environments excluded *Anthracoceras* from the northern part of the area, but it continued to live within a zone approximately 40 km wide at the shelf margin (figs. 5 and 8) in southeastern Nevada and northeastern Arizona (table 1). The occurrence of the genus approximately 100 km landward from the shelf margin in extreme southeastern Nevada (locality 1) may be the result of incorrect location of the shelf margin due to post-Permian or to post-Laramide structural extension in this structurally complex area.

In the Lower Moore *Anthracoceras* became extinct by the end of the early middle Visean (Composite Zone 14), although seemingly favorable shelf carbonate environments continued into Composite Zone 15 on the Leadville Redwall shelf (localities 1–3 on fig. 8). No pre-Pogonip occurrences are known.

Western Canada

In Canada, *Anthracoceras* has been found in a much more restricted area than in the United States. It is known only from east-central British Columbia (figs. 1 and 9), where it occurs in 10 closely spaced outcrop sections within the overthrust belt of the Rocky Mountain fold-and-thrust belt (fig. 9). Most of the occurrences are in a narrow belt of protected-shelf carbonate deposited slightly landward of the shelf margin of a carbonate platform (figs. 9–11) (see appendix C for complete descriptions of localities). The protected-shelf deposits extend northeastward from Jarvis Lakes in southeast Mackenzie Pass map area (NTS 93I) into the Yahatinda River region of the Pine Pass map area (NTS 93O). Numerous specimens have also been collected from middle-slope carbonates (fig. 11) deposited southwest of the shelf-margin positions both at the northeastern part of the study area (figs. 9 and 10, localities 1 and 2). No specimens of *Anthracoceras* have been found elsewhere in the Canadian Cordillera or in the Interior platform to the east, although more than 100 coral collections from correlative strata have been examined. The belt of protected-shelf, shelf-margin, and slope carbonates extends northeastward from the study area into the subsurface (figs. 3 and 9), where very few coral collections are available. South of the study area, the genus has not been observed in the numerous occurrences of protected-shelf and shelf-margin facies examined, but in this area most of these facies have not been thoroughly sampled. Also, few coral collections have been made from this stratigraphic interval from slope carbonate rocks in areas other than the study area (fig. 3). Therefore, the apparent geographic restriction of *Anthracoceras* in Canada is probably an artifact of sampling, but the possibility that it is related to undetermined environmental factors cannot be ruled out.

Anthracoceras has the same age range (Composite Zones 12 to 14) in Canada as in the United States (fig. 12). Localities 5 and 9 have yielded specimens from the basal part of Composite Zone 12, but most occurrences are in Composite Zone 13 or in the lower part of Composite Zone 14. The vertical distribution of these occurrences shows no obvious geographic pattern. Their zonal assignments are based on biostratigraphic data from associated brachiopod faunas. The assignment of specimens from locality 2 was estimated from their stratigraphic position by correlation with nearby well-dated sections.

Anthracoceras lived on the protected shelf and middle slope in a genetic differentiated mobile Visean carbonate platform that had a broad, sand-dominated shelf margin and low-gradient slope (fig. 11). The main lithofacies belts of the platform are incompletely preserved in the Turner Valley Formation, the lower part of the Mount Head Formation, and the Prophet Formation (fig. 10). The regional and local lithostratigraphic, lithofacies, and depositional environments of these formations have been discussed by Beauchamp and others (1989), Mamet and others (1990), and Richards and others (in press). A complete shelf-to-basin transition is preserved only in a series of sections (localities 1–3, fig. 9) in the northeastern part of the area. This transition records a westward progradation of the protected shelf, shelf margin, and slope. Elsewhere in the study area, protected-shelf deposits containing *Anthracoceras* taxon are widely preserved, but shelf-margin deposits are preserved only at locality 10 (fig. 9), and slope lithofacies are absent. In east-central British Columbia, the protected-shelf deposits containing *A. taylori* (figs. 9 and 10) formed near the seaward limit of the protected shelf and grade basinward (southwestward) into coeval shelf-margin and upper slope facies. The protected-shelf deposits are mainly peloid- and pellet-rich bedded, mixed oncolital grainstone and packstone deposited in shallow subtidal (above storm wave base) environments (fig. 11). These deposits generally occur in the Turner Valley Formation and Banff(?) Member of the Mount Head Formation, but some of them

Figure 9. Present supratidaspastic facies distribution of Vocan (Composite Zones 12 to 14) deposits, east-central British Columbia (see study area, fig. 2) and Anshelasma occurrences in relation to shelf margin.

Figure 10. Stratigraphic cross section A–A' showing Anshelasma occurrences (appendix C, localities 1–30) and depositional environments (fig. 11) in Kindle Group and Besa River Formation, study area, east-central British Columbia. Line of cross section shown on figure 9. Adapted from Richards and others (in press).

are in the Wilemanic Member of the Mount Head Formation (figs. 10 and 12). Toward the east, the broad belt of protected-shelf lithofacies commonly records deepening. Farther eastward, protected-shelf lithofacies indicate shallowing, and eventually grade into intertidal to supratidal carbonates and anhydrite deposited on the inner protected shelf and on a vast restricted shelf (figs. 10 and 11).

The shelf-margin and upper-slope facies are mainly bryozoan-peloid-foram grainstone (fig. 11). In the study area, this belt of high-energy deposits, which has not been identified near Anshelasma localities in the United States, is thin and relatively narrow compared to that of southwestern Alberta and southeastern British Columbia. In the latter region, the belt of shelf-margin grainstone is more than 30 km wide.

Toward the northwest, the shelf-margin and upper-slope grainstone grades into finer grained middle-slope and upper-slope lithofacies that occur in the Prophet Formation (figs. 9 and 10; localities 1 and 2) and locally contain abundant Anshelasma. The coral-bearing slope deposits are bioturbated, cherty, peloid-foram-spicule packstone. Corals in these deposits are very well preserved but appear to have been reworked, because they are not in growth position. They are not, however, associated with either shallow-water fossils or abundant bryozoans and peloid-foram oncolites, the principal allochems of upper-slope and shelf-margin deposits. We conclude that these corals lived on the middle slope and did not originate on the upper slope or shelf margin. These middle-slope dwellers (fig. 11) are represented by a new species of *Anshelasma* (fig. 2 *E*, *F*) known from no other localities in Canada or the United States. The unique occurrence of this species in Composite



Figure 11. Conventional sedimentological model of Cretaceous turbidite platform (from Richards and others, in press).

Zone 14 at these localities may represent a migration of the genus down the slope from its earlier habitat behind the shelf margin toward the end of the progression (fig. 9).

The distribution of *Inoceramus* suggests that the genus has considerable utility as a paleoenvironmental indicator. In post-Cretaceous British Columbia, this readily identifiable and commonly abundant extinct fossil principally in neritic settings that were near or above storm wave base and were located on either side of the shelf margin grain-stone belt (fig. 11). All identified specimens of *A. typicus* came from deposits of the outer protected shelf, thereby suggesting that the species lived mainly on the outer protected shelf. A. sp. aff. from localities 1 and 2 may be a slope indicator because it has been identified only from middle-slope deposits. *Inoceramus* has not been observed in deep-water, shelf-margin (= intertidal) deposits, and where it occurs in protected-shelf facies, these neritic deposits formed near the shelf margin. The fact that *Inoceramus* has not been found east of the broad shelf-margin grainstone belt in Canada (table 3) the study area may be related to the greater degree of restriction of the protected shelf in that area.

INFERRED SHELF-MARGIN LOCATIONS

Western United States

In northwestern Montana, western Alberta and eastern Mesozoic and Cenozoic uplift removed most of the record of Upper deposition (Klove and Lantz, 1948; Mudge and others, 1962; Hunter, 1977). Hence the location of the shelf margin is unknown. The occurrences of *Inoceramus* at localities 12 and 13 provide a basis for estimating the shelf margin location (fig. 10). By using the *Inoceramus*



Figure 13. Chronometric lithofacies profiles from *Ankhelasma* localities 12 and 13 in westernmost Carboniferous outcrops in northwestern Montana to inferred shelf margin. Location of control points shown in figs. 6 and 7 and control points described in appendix D. Lost record indicated by shaded pattern. See figure 8 for explanation of other symbols.

distribution model (see p. B4–B11). These occurrences at localities 12 and 13 are in the lower half of Composite Zone 13 (fig. 5).

The *Ankhelasma* occurrence at Little Blackfoot River (locality 12, fig. 6) in profile *G–G'* (fig. 13) suggests a shelf-margin position no more than 50 km to the west of the *Ankhelasma* location, using the distribution model based on known distance from the shelf margin (table 1). However, outcrops of Viséan rocks between locality 12 and Rattler Gulch, 65 km to the west, do not reveal a basinal facies in Composite Zone 13—suggesting a shelf-margin position west of Rattler Gulch. The present position of the inferred shelf margin in this profile (*G–G'*) is based on westward projection of lithofacies trends; it is approximately 75 km west of locality 12 (fig. 13). This anomaly is thought to be the result of post-Laramide structural extension along this profile, on the basis of recent recognition of structural extension in the thrust belt (W.J. Perry, oral commun., 1987).

The *Ankhelasma* occurrence at Hannan Gulch (locality 13, fig. 6) in profile *H–H'* (fig. 13) is compatible with the distribution model. Westward projection of lithofacies trends in this profile indicates a shelf-margin position 45 km west of the *Ankhelasma* location (fig. 13).

Western Canada

In the overthrust belt of east-central British Columbia, the protected-shelf occurrences of *Ankhelasma* are located in a narrow, northwest-trending outcrop belt subparallel to the depositional and structural strike of Viséan rocks (figs. 3, 9, and 10). Viséan shelf-margin deposits are preserved only in the northwestern and southeastern parts of the Carboniferous outcrop belt. Most *Ankhelasma* occurrences on the protected shelf lie east of an area of lost Viséan record. The relative present position of the shelf-margin deposits is known for only two of these occurrences (localities 9 and 10; figs. 9 and 10). The eastern limits of the shelf margin deposits assigned to Composite Zones 12 to 14 were projected into the area of lost record by means of lithofacies analyses based on unpublished field observations. The placement of these projected limits is supported by comparison with the occurrences of *Ankhelasma* shown on figure 9. All protected-shelf occurrences of the genus lie within 11 km of the known and projected eastern shelf-margin limits, well within the present-day width of the *Ankhelasma* belt established for the Western United States.

The maximum present width of the main *Ankhelasma* belt on the protected shelf in Canada is less than the average width (about 22 km) of the *Ankhelasma* belt in the United States (table 1). This may be only an apparent difference resulting from different definitions of the shelf margin in the two areas. In Canada, the shelf margin is defined by a grainstone belt that is more than 10 km wide (fig. 9), whereas in the United States, the margin is defined by an arbitrary line separating shelf and off-shelf lithofacies (figs. 6 and 7). Laramide thrusting and folding also may have reduced the widths of the Carboniferous facies belts more in Canada than in the United States.

CONCLUSIONS

(1) The distinctive, endemic rugose coral genus *Ankhelasma* Sando, 1961, occurs only in a narrow geographic belt in rocks of early and middle Viséan age in the Western Interior region of North America. With very few exceptions, this *Ankhelasma* belt is located on the landward side of the Viséan shelf margins.

(2) Only two species have been recognized thus far in all the known occurrences of the genus. *Ankhelasma typicum* Sando, 1961, the more abundant species, lived exclusively in shallow, subtidal environments of the outer protected shelf within approximately 50 km of the shelf margin during the early and middle Viséan. Toward the end of the phylogenetic life span of *Ankhelasma*, an undescribed new species lived on the middle slope of a middle Viséan carbonate platform on the seaward side of the shelf margin in east-central British Columbia (Peace River embayment).

(3) The model of *Ankhelasma* distribution, based on known occurrences, can be used to predict locations of Viséan shelf margins in areas where the Viséan depositional record was removed by erosion or obscured by later tectonism. This model may also be used to confirm or deny shelf margin projections by lithofacies analyses into areas where lithofacies data are sparse or unavailable.

REFERENCES CITED

Beauchamp, B., Richards, B.C., Bamber, E.W., and Mamet, B.L., 1980, Lower Carboniferous lithostratigraphy and carbonate facies, upper Banff Formation and Rundle Group, east-central British Columbia, *in* Current Research, pt. A: Geological Survey of Canada Paper 80-1A, p. 627–644.

Burbiggen, E.T., 1964, Geology of the Black Mountain area, Lewis and Clark and Powell Counties, Montana: Princeton, New Jersey, Princeton University, Ph.D. thesis, 120 p.

Craig, L.C., and Connor, C.W., coordinators, 1979, Paleotectonic investigations of the Mississippian System in the United States: U.S. Geological Survey Professional Paper 1010, pt. III, 35 plates.

Dockal, J.A., 1980, Petrology and sedimentary facies of Redwall Limestone (Mississippian) of Uinta Mountains, Utah and Colorado: Iowa City, Iowa, University of Iowa, Ph.D. thesis, 423 p.

Gordey, S.P., Abbott, J.G., Templeman-Kluit, D.J., and Gabrielse, H., 1987, "Antler" clastics in the Canadian Cordillera: Geology, v. 15, no. 2, p. 103–107.

Gutschick, R.C., and Sandberg, C.A., 1983, Mississippian continental margins of the conterminous United States: Society of Economic Paleontologists and Mineralogists Special Publication 33, p. 79–96.

Gutschick, R.C., Sandberg, C.A., and Sando, W.J., 1980, Mississippian shelf margin and carbonate platform from Montana to Nevada, *in* Fouch, T.D., and Magathan, E.R., eds., Paleozoic paleogeography of west-central United States: Rocky Mountain Section, Society of Economic Paleontologists and Mineralogists, West-Central Unites States Paleogeography Symposium 1, p. 111–128.

Hamm, T.E., 1977, Lower Mississippian sedimentation in southwestern Montana: Rolla, Missouri, University of Missouri-Rolla, Ph.D. thesis, 116 p.

Hall, O.K., 1967, The Mississippian System across the Wasatch line, east-central Idaho, extreme southwestern Montana: Montana Geological Society, Eighteenth Annual Field Conference Guidebook, p. 51–62.

———1968, Mississippian stratigraphy and sedimentation across the Wasatch line, east-central Idaho and extreme southwestern Montana: State College, Pennsylvania, Pennsylvania State University, Ph.D. thesis, 178 p.

Lane, H.R., Sandberg, C.A., and Ziegler, W., 1980, Taxonomy and phylogeny of some Lower Carboniferous conodonts and preliminary standard post-*typomeandra* zonation: Geologica et Palaeontologica, v. 14, p. 117–164.

Langenheim, R.L., Jr., 1962, Nomenclature of the Late Mississippian White Pine Shale and associated rocks in Nevada: Illinois State Academy of Science Transactions, v. 55, no. 2, p. 133–145.

———1963, Mississippian stratigraphy in southwestern Utah and adjacent parts of Nevada and Arizona: Intermountain Association of Petroleum Geologists, Twelfth Annual Field Conference Guidebook, p. 30–41.

Mamet, B.L., and Skipp, Betty, 1970, Lower Carboniferous calcareous foraminifera, preliminary zonation and stratigraphic implications for the Mississippian of North America: International Congress of Carboniferous Stratigraphy and Geology, 6th, Sheffield, 1967, Comptes Rendus, v. 3, p. 1129–1146.

Mamet, B.L., Bamber, E.W., and Macqueen, R.W., 1986, Microfacies of an Upper Carboniferous Banff Formation and Rundle Group, Monkman Pass map area, northeastern British Columbia: Geological Survey of Canada Bulletin 353, 91 p.

Madge, M.R., 1972, Pre-Quaternary rocks of the Sun River Canyon area, northwestern Montana: U.S. Geological Survey Professional Paper 663-A, 142 p.

Madge, M.R., Sando, W.J., and Dutro, J.T., Jr., 1962, Mississippian rocks of the Sun River Canyon area, Sawtooth Range, Montana: American Association of Petroleum Geologists Bulletin, v. 46, no. 11, p. 2003–2018.

Nichols, K.M., 1986, Regional significance of lithologic correlation of Mississippian rocks at Pentagon Mountain and the Sawtooth Range, northwestern Montana: U.S. Geological Survey Open-File Report 86-39, 1 sheet.

Nilsen, T.H., 1977, Paleogeography of Mississippian turbidites in south-central Idaho, *in* Stewart, J.H., Stevens, C.H., and Fritsche, A.E., eds., Paleozoic paleogeography of the Western United States: Pacific Section, Society of Economic Paleontologists and Mineralogists, Pacific Coast Paleogeography Symposium 1, p. 175–209.

Parker, J.W., and Roberts, J.W., 1963, Devonian and Mississippian stratigraphy of the central part of the Colorado Plateau: Four Corners Geological Society, Fourth Field Conference, Symposium on Shelf Carbonates of the Paradox Basin, p. 31–60.

Pierce, R.W., and Langenheim, R.L., Jr., 1974, Platform conodonts of the Monte Cristo Group, Mississippian, Arrow Canyon Range, Clark County, Nevada: Journal of Paleontology, v. 48, no. 1, p. 149–169.

Poole, F.G., 1974, Flysch deposits of the Antler foreland basin, Western United States, *in* Dickinson, W.R., ed., Tectonics and sedimentation: Society of Economic Paleontologists and Mineralogists Special Publication 22, p. 58–82.

Poole, F.G., and Sandberg, C.A., 1977, Mississippian paleogeography and tectonics of the Western United States, *in* Stewart, J.H., Stevens, C.H., and Fritsch, A.E., eds., Paleozoic paleogeography of the Western United States: Pacific Section, Society of Economic Paleontologists and Mineralogists, p. 67–85.

Richards, B.C., Bamber, E.W., Higgins, A.C., and Utting, J., *in* press, Carboniferous, *in* Stott, D.F., and Aitken, J.D., eds., Sedimentary cover of the craton: chap. 4E (Stratigraphy): The geology of North America: Geological Survey of Canada, v. 5.

Rose, P.R., 1976, Mississippian carbonate shelf margins, Western United States: U.S. Geological Survey Journal of Research, v. 4, no. 4, p. 449–466.

Sandberg, C.A., and Gutschick, R.C., 1984, Distribution, microfauna, and source rock potential of Mississippian Delle Phosphatic Member of Woodman Formation and equivalents, Utah and adjacent states, *in* Woodward, J., Meissner, F.F., and Clayton, J.L., eds., Hydrocarbon source rocks of the Greater Rocky Mountain region: Rocky Mountain Association of Geologists, p. 135–178.

Sandberg, C.A., Gutschick, R.C., Johnson, J.G., Poole, F.G., and Sando, W.J., 1982, Middle Devonian to Late Mississippian geologic history of the Utah-Idaho trough and overthrust belt region, Western United States—A summary: Rocky Mountain Association of Geologists–Geologic Studies of the Cordilleran Thrust Belt, v. 2, p. 691–719.

Sando, W.J., 1961, Morphology and ontogeny of Ankhelasma, a new Mississippian coral genus: Journal of Paleontology, v. 35, no. 1, p. 65–81.

———1976, Mississippian history of the northern Rocky Mountains region: U.S. Geological Survey Journal of Research, v. 4, no. 3, p. 317–338.

———1985, Revised Mississippian time scale, Western Interior region, conterminous United States: U.S. Geological Survey Bulletin 1605-A, p. A15–A26.

———1988, Madison Limestone (Mississippian) paleokarst—A geologic synthesis, *in* James, N.P., and Choquette, P.W., eds., Paleokarst: New York, Springer-Verlag, p. 256–277.

Sando, W.J., and Bamber, E.W., 1985, Coral zonation of the Mississippian System in the Western Interior province of North America: U.S. Geological Survey Professional Paper 1334, 81 p.

Sando, W.J., Bamber, E.W., and Armstrong, A.K., 1975, Endemism and similarity indices: Clues to the zoogeography of North American Mississippian corals: Geology, v. 3, no. 11, p. 661–664.

———1977, The zoogeography of North American Mississippian corals: Second International Symposium on Corals and Fossil Coral Reefs, Bureau de Recherches Geologiques et Minières, Memoire 89, p. 175–184.

Sando, W.J., and Dutro, J.T., Jr., 1981, Some stratigraphic sections of the Madison Group in the overthrust belt of western Wyoming and southeastern Idaho: U.S. Geological Survey Open-File Report 81-1354, 39 p.

Sando, W.J., Dutro, J.T., Jr., and Gere, W.C., 1959, Brazer Dolomite (Mississippian), Randolph quadrangle, northeast Utah: American Association of Petroleum Geologists Bulletin, v. 43, no. 12, p. 2741–2766.

Sando, W.J., Sandberg, C.A., and Gutschick, R.C., 1981, Stratigraphic and economic significance of Mississippian sequence at North Georgetown Canyon, Idaho: American Association of Petroleum Geologists Bulletin, v. 65, no. 8, p. 1435–1443.

Sando, W.J., Sandberg, C.A., and Perry, W.J., Jr., 1985, Revision of Mississippian stratigraphy, Tendoy Mountains, southwest Montana: U.S. Geological Survey Bulletin 1656-A, 10 p.

Skipp, Betty, Hammer, I.R., and Dertz, D.E., 1984, Geology, geochemistry, and mineral resource potential of the Eighteen

mile Wilderness Study Area (ID-43-3), Lemhi County, Idaho: U.S. Geological Survey Open-File Report 84-279, 55 p.

Sloss, L.L., and Laird, W.M., 1945, Mississippian and Devonian stratigraphy of northwestern Montana: U.S. Geological Survey Oil and Gas Preliminary Chart 15.

Sproule, J.C., 1958, Petroleum and natural gas prospects of Mississippian and Jurassic of western Canada, *in* Goodman, A.J., ed., Jurassic and Carboniferous of western Canada: American Association of Petroleum Geologists, p. 3–9.

Steed, D.A., 1980, Geology of the Virgin River Gorge, northwest Arizona: Brigham Young University Geology Studies, v. 27, pt. 3, p. 99–115.

Struik, L.C., 1987, The ancient western North American margin: An alpine rift model for the east-central Canadian Cordillera: Geological Survey of Canada Paper 87-15, 19 p.

Templeman-Kluit, D.J., 1979, Transported cataclasite, ophiolite and granodiorite in Yukon: Evidence of arc-continent collision: Geological Survey of Canada Paper 79-14, 27 p.

APPENDIX A. *ANKHELASMA* LOCALITIES IN THE UNITED STATES

1. Potosi Mine

Location: CN½ SW¼ NE¼ sec. 12, T. 23 S., R. 57 E., Shenandoah Peak quadrangle (15 min), Goodsprings district, Clark County, Nevada.

Stratigraphic position: Yellowpine Member of Monte Cristo Limestone, 0–1.2 m (0–4 ft) above base.

Collections: • USGS 27968–PC, 0.9–1.2 m (3–4 ft) above base. Approximately 150 silicified specimens of *A. typicum* Sando etched from limestone.

• Sandberg Sample PO-3, lower 0.09 m (0.3 ft). Two silicified specimens of *A.* sp. indet. etched from limestone.

Lithology: Thick-bedded, fine- to medium-grained crinoidal limestone. Collected sample is dark, crinoidal packstone.

Age: Associated fossils (USGS 27968–PC) are *Canadiphyllum?* sp., *Syringopora* sp., and Foraminifer Zone 12 microfossils (table A1). The occurrence is regarded as high in Composite Zone 14 because the underlying Arrowhead Member contains Foraminifer Zone 12 microfossils (table A1). (USGS 27967–PC), and overlying beds in the Yellowpine Member yielded Foraminifer Zone 12/13 microfossils (table A1) (USGS 27969–PC).

2. Tungsten Gap (see Pierce and Langenheim, 1974, for published stratigraphic section).

Location: CN½ sec. 10, T. 15 S., R. 64 E., Arrow Canyon quadrangle (15 min), Arrow Canyon Range, Clark County, Nevada.

Stratigraphic position: Yellowpine Member of Monte Cristo Limestone, 1.5–7.6 m (5–25 ft) above base.

Collections: • USGS 27974–PC, 7.6 m (25 ft) above base. Approximately 40 calcareous specimens of *A. typicum* Sando in limestone.

• USGS 27841–PC, lower 6.1 m (20 ft). Approximately 20 calcareous specimens of *A. typicum* Sando in limestone.

• USGS 27842–PC, 1.5–2.4 m (5–8 ft) above base. Approximately 100 calcareous specimens of *A. typicum* Sando in limestone.

Lithology: Thick-bedded, dark, fine- to medium-grained crinoidal limestone. Collected samples are dark, medium-grained crinoidal grainstone.

Age: Foraminifer control is lacking in this section. The occurrence of *Siphonodendron* sp., *Vesiculophyllum* sp., *Amplexizaphrentis* sp., and *Syringopora* sp. suggests Coral Zone IIIA or B. Conodont evidence suggests that the *Ankhelasma* occurrence is high in Composite Zone 14, similar to the occurrence at Potosi Mine (C.A. Sandberg, oral commun., 1986).

3. Virgin River Gorge (see Steed, 1980, for general geology of this locality).

Location: Secs. 20 and 30, T. 41 N., R. 14 W., Littlefield quadrangle (15 min), Mohave County, Arizona.

Stratigraphic position: Lowermost part of Yellowpine Member of Monte Cristo Limestone.

Age: Abundant specimens of *Ankhelasma* sp. indet. were identified in the field by C.A. Sandberg (oral commun., 1986). No fossils were collected, but this occurrence is presumably the same stratigraphic level and age (Composite Zone 14) as occurrences at Potosi Mine and Tungsten Gap.

4. Samak (see Dockal, 1980, for additional data).

Location: S½ NE¼ NE¼ sec. 25, T. 2 S., R. 6 E., Woodland quadrangle (7.5 min), Uinta Mountains, Summit County, Utah.

Stratigraphic position: Brazer Dolomite, 1.5 m (5 ft) below top of Mission Canyon equivalent (member 2 of the Brazer).

Collection: • USGS 27065–PC. One silicified specimen of *A.* cf. *A. typicum* Sando etched from limestone.

Lithology: Medium-bedded, dark, fine- to medium-grained crinoidal limestone. Collected sample is partly dolomitized crinoidal packstone.

Age: *Ankhelasma* occurs here with *Siphonodendron ocalinum* Sando, *Sychnoelasma* sp., *Vesiculophyllum* sp., *Lophophyllum?* sp., *Amplexizaphrentis* sp., and

Table A1. Foraminifers and algae in samples from the United States cited in appendix A

[All samples are registered in U.S. Geological Survey Upper Paleozoic Catalog, except for BSH-2A, which is in the catalog of C. A. Sandberg. Imp./Sciences of Fossils by R.C. Moore]

Taxa	Robison Mine			Trail Hollow	Emma Canyon	Unnamed Canfield Mb.	Hot Dog Peak	Sheep Creek			Big Sheep Creek		Bell Canyon		Little Blackfoot River
	27065	27066	27067	27074	16936	16937	17006	20222	20223	20227	BSH-2A	20665	29401	29409	21401
Attenuloplica sp.			X												
Bravoia sp.	X														
Calcifolium? lutrix Wilkerson					X		X	X			X			X	
C. pachytheca (Prosser)				X	X	X	X	X	X	X	X	X			
C. sp.	X	X	X									X			X
Dielasma(?) *D. arcuatum* Mansuy			X												
D. sp.			X												
Eurlasmella crassula (Beecher)									X						
E. calgaryi (Kindle Cheiloniformis and Rothpletz)		X	X	X	X			X	X			X	X	X	X
E. sp.						X	X	X		X					
Ghaniini sp.		X										X			
Lingulonis sp.	X	X	X	X		X						X	X	X	X
cf. *Endothyranopsis*? sp.	X														
Endothyranopsis bilaterica (Moller)		X	X											X	
E. volvida (Toomey)	X														
E. spinosa (Moller)	X		X		X										
E. spinosa spinosa (Zeller)		X							X						
E. of the group *E. spinosa* (Zeller)								X	X			X		X	X
E. sp.								X					X		
Eolasmoides moelleri (Malakhova in Dain)		X													
E. sp.	X	X	X									X		X	X
Eoplectodella media Vdovenko				X	X		X	X							
Geinitzella sp.	X														
Globoendothyra bogdanovi (Skipp)		X													
G. of the group *G. baileyi* (Hall)				X				X	X		X	X			X
G. of the group *G. sinuosa* (Grozdilova in Lebedeva)	X													X	
G. sp.			X		X		X		X	X				X	
G.? *visensis* Zeller											X				
Ivanella davisensis Rettinger												X			
I. sp.											X				
Koskinobigenina adkinsi (de Kornfeld)														X	
K. sp.		X													
Loeblichella sp.					X	X				X	X				
Mametella sp.		X							X				X		
Orthocoeliopsis sp.							X			X					
Orthovirgulina sp.		X													X
Ozawainella sp.										X				X	
Palaeotextularia lahuseni (von Moller)												X			
P. sp.							X								X
Pseudoammovertella sp.								X			X	X			
Pseudoendothyra sp.															X
Pseudotaxis sp.	X		X	X	X	X	X	X	X		X	X	X	X	X
Pseudovidalina sp.														X	X
Septatournayella sp.										X					
S. sp.	X	X	X										X		
cf. *Stigmella*? sp.					X			X							
cf. *Spinoendothyra*? sp.											X				
Stacheiidae									X						
Stacheoides sp.	?														X
Tetrataxis sp.	X			X					X						X

Syringopora morphogroup C (USGS 27065–PC), which are indicative of Coral Zone IIIA (near base). Conodonts from this stratigraphic level represent either the *Scaliognathus anchoralis-Doliognathus latus* Zone or the *Gnathodus texanus* Zone (C. A. Sandberg, oral commun., 1978). These fossils suggest a zonal position at or near the base of Composite Zone 12.

5. Trail Hollow (see Dockal, 1980, for additional data).

Location: NW¼ sec. 33, T. 3 S., R. 9 W., Iron Mountain quadrangle (7.5 min), Uinta Mountains, Wasatch County, Utah.

Stratigraphic position: Brazer Dolomite, 1.5–7.6 m (5–25 ft) below top of Mission Canyon equivalent (member 2 of the Brazer).

Collections: ● USGS 27074–PC. Approximately 100 calcareous specimens of *A. typicum* Sando in limestone.

Lithology: Thin- to thick-bedded, weakly laminated, coarse-grained bioclastic limestone. Collected sample is dark, partly dolomitized coral packstone.

Age: This sample contains the same assemblage of corals and conodonts as USGS 27065–PC at Samak. It also contains foraminifers that have a possible range of Foraminifer Zone 10 to 12 (table A1). A zonal range at or near the base of Composite Zone 12 is indicated.

6. Crawford Mountains (see Sando and others, 1959, and Sando, 1961, for additional data). All specimens belong to *Ankhelasma typicum* Sando and are from the Randolph quadrangle (15 min), Rich County, Utah.

A. Unnamed locality.

Location: NW¼ NW¼ sec. 29, T. 11 N., R. 8 E.

Stratigraphic position: Brazer Dolomite, upper member (member 3), exact position unknown.

Collection: ● USGS 486–PC. Three specimens preserved as molds and casts in chert.

Lithology: Collected sample is cherty, dark, fine-grained calcitic dolomite.

Age: The sample contains nondiagnostic brachiopods and corals. It is dated as middle Composite Zone 12 on the basis of similarity of its stratigraphic position to collections from nearby localities that have been dated by diagnostic fossils.

B. Brazer Canyon (type section of Brazer Dolomite).

Location: SE¼ NW¼ sec. 20, T. 11 N., R. 8 E.

Stratigraphic position: Brazer Dolomite, 44.5–50.6 m (146–166 ft) above base of upper member.

Collections: ● USGS 16906–PC, 44.5 m (146 ft) above base. Two poorly preserved calcitic and dolomitic specimens in dolomite.

● USGS 16907–PC, 48.2 m (158 ft) above base. Two poorly preserved calcitic and dolomitic specimens in dolomite.

● USGS 16908–PC, float from 41.2–50.6 m (135–166 ft) above base. Eleven poorly preserved calcitic and dolomitic specimens in dolomite.

Age: These samples contain *Zaphriphyllum* sp., *Camaliphyllum* sp., and *Vesiculophyllum*? sp., which suggest Coral Zone IIIA. A zonal position near the middle of Composite Zone 12 is suggested by proximity to Coral Zone II corals in the underlying Mission Canyon equivalent (member 2 of the Brazer). No conodonts or foraminifers were found.

C. Emma Canyon.

Location: C sec. 17, T. 11 N., R. 8 E.

Stratigraphic position: Brazer Dolomite, 44.2–56.4 m (145–185 ft) above base of upper member (member 3).

Collections: ● USGS 16936–PC, 48.2 m (156 ft) above base. Approximately 50 specimens preserved as molds and casts in chert.

● USGS 16938–PC, 56.4 m (185 ft) above base. Six poorly preserved calcitic and dolomitic specimens in dolomite.

● USGS 16939–PC, 44.2 m (145 ft) above base. Three poorly preserved calcitic and dolomitic specimens in dolomite.

Lithology: Same as locality 6B; chert preserves pelletal packstone texture.

Age: The interval containing *Ankhelasma* at this locality is approximately the same as that at locality 6B, which has been dated as middle Composite Zone 12. USGS 16936–PC at Emma Canyon contains microfossils of Foraminifer Zone 10 or 11 (table A1), which confirms the Composite Zone 12 determination.

D. Rex Peak.

Location: N½ sec. 31, T. 11 N., R. 8 E.

Stratigraphic position: Brazer Dolomite, 37.5–40.5 m (123–133 ft) above base of upper member (member 3).

Collections: ● USGS 17717-PC, exact position unknown. Several hundred poorly preserved silicified specimens etched from dolomite.

● USGS 17722-PC, 37.5–40.5 m (123–133 ft) above base. 285 well-preserved silicified specimens and several hundred poorly preserved ones etched from dolomite.

Lithology: Same as locality 6B.

Age: Same as locality 6B.

E. Unnamed Crawford Mountains

Location: NW¼ NW¼ sec. 6, T. 10 N., R. 8 E.

Stratigraphic position: Brazer Dolomite, upper member (member 3); exact position unknown.

Collection: ● USGS 18074-PC, exact position unknown. Approximately five poorly preserved calcitic and dolomitic specimens in dolomite.

Lithology: Same as locality 6B.

Age: Same as locality 6B. USGS 18074-PC contains microfossils of Foraminifer Zone 10 or 11 (table A1), which confirms the Composite Zone 12 determination for *Ankhelasma* at this locality.

7. Haystack Peak (see Sando and Dutro, 1981, for description of stratigraphic section).

Location: NW¼ NE¼ sec. 27, T. 34 N., R. 117 W., Bedford quadrangle (7.5 min), Salt River Range, Lincoln County, Wyoming.

Stratigraphic position: Mission Canyon Limestone, 94.5–97.5 m (310–320 ft) below top (unit 49 of Sando and Dutro, 1981, p. 12).

Collection: ● USGS 17900-PC. One partly silicified calcitic specimen in limestone and two silicified specimens etched from limestone, all identified as *A.* cf. *A. typicum* Sando.

Lithology: Thin- to medium-bedded, medium- to very coarse grained crinoidal bioclastic limestone. Collected sample is dark crinoidal packstone.

Age: This occurrence is well documented with corals of Zone IIIA (*Siphonodendron ocalinum* Sando; *Zaphriphyllum* sp., *Amplexizaphrentis* sp., *Vesiculophyllum* sp., *Syringopora* sp., *Michelinia* morphogroups B and C; *Pleurophonella* morphogroup A) that occur in the same bed (USGS 17900-PC) and above (USGS 17901-PC), and below (USGS 17899-PC) a Foraminifer Zone 9/10 microfossils occur below the collection (table A1) (USGS 17906-PC). A zonal position near the middle of Composite Zone 12 is indicated for *Ankhelasma* here.

8. Sheep Creek (see Sando and Dutro, 1981, for description of stratigraphic section).

Location: E½ sec. 28, T. 1 N., R. 45 E., Irwin quadrangle (30 min), Snake River Range, Bonneville County, Idaho.

Stratigraphic position: Mission Canyon Limestone, 93–96 m (305–315 ft) below top (unit 56 of Sando and Dutro, 1981, p. 86).

Collection: ● USGS 20122-PC. One calcitic specimen of *A.* cf. *A. typicum* Sando in limestone.

Lithology: Medium- to thin-bedded, dark, predominantly fine-grained limestone. Collected sample is dark, crinoidal, foraminiferal packstone.

Age: This occurrence is well documented with corals of Zone IIIA (*Siphonodendron ocalinum* Sando; *Zaphriphyllum* sp., *Amplexizaphrentis* sp., *Vesiculophyllum* sp., *Syringopora* morphogroups B and C). The corals occur with microfossils of Foraminifer Zones 10–11 (table A1) (USGS 20122-PC) and occur below microfossils of Foraminifer Zone 11 (table A1) (USGS 20121-PC) and above microfossils of Foraminifer Zone 10(?) (table A1) (USGS 20117-PC). A zonal position in the upper half of Composite Zone 12 is indicated for *Ankhelasma* here.

9. Chamberlain Creek (see Skipp and others, 1984, pl. 1, for geologic map of the area).

Location: SE¼ sec. 20, T. 14 N., R. 29 E., Morrison Lake quadrangle (15 min), Beaverhead Mountains, Lemhi County, Idaho.

Stratigraphic position: Debris flow in Middle Canyon Formation, exact position unknown.

Collection: ● USGS 29266-PC. One calcitic specimen of *A.* cf. *A. typicum* Sando in limestone.

Lithology: Collected sample is dark, silty and argillaceous, phosphatic, coarse-grained crinoidal grainstone.

Age: Other megafossils collected from the locality (USGS 29584-PC) include indeterminate horn corals, euomphalacean gastropods, a spiriferoid brachiopod, and an orthoceracone cephalopod; none of which is diagnostic for age determination. Matrix from USGS 29584-PC yielded zonally indeterminate conodonts *Bispathodus aculeatus* and *Polygnathus communis communis*, which have a range from Osagean to middle Meramecian (late Tournaisian to middle Visean) (C.A. Sandberg, written commun., 1985). The deposit is interpreted as a debris flow derived from shallow-water-shelf sediments of Composite Zone 12 age (Mission Canyon Limestone) and redeposited in a deep basin represented by the Middle Canyon Formation. Dating is based on the age of *in situ* occurrence at Big Sheep Creek (locality 10).

10. Big Sheep Creek.

Location: SW¼ NE¼ sec. 22, T. 14 S., R. 10 W., Gallagher Gulch quadrangle (7.5 min), Tendoy Mountains, Beaverhead County, Montana.

Stratigraphic position: Mission Canyon Limestone, upper member, approximately 6.7 m (22 ft) above base.

Collection: ● USGS 28605-PC. Six calcitic specimens of *A.* cf. *A. typicum* Sando in limestone.

Lithology: Collected sample is dark, partly dolomitized crinoidal packstone.

Age: Corals of Zone IIIA (*Vesiculophyllum* sp., *Siphonodendron ocalinum* Sando, and *Syringopora* sp.) occur immediately above the bed containing *Ankhelasma*. Microfossils of Foraminifer Zone 10–11 (table A1) (Sandberg Sample BSH–2A) occur at the same level as the collected sample. A zonal position near the base of Composite Zone 12 is indicated for *Ankhelasma* here.

11. Bell Canyon (see Sando and others, 1985, for section description).

Location: NW¼ NE¼ SW¼ sec. 17, T. 11 S., R. 10 W., Red Rock and Kidd quadrangles (7.5 min), Tendoy Mountains, Beaverhead County, Montana.

Stratigraphic position: Mission Canyon Limestone, near top of lower(?) member to 26 m (85.3 ft) below top of upper member.

Collections: ● USGS 29305-PC, 26 m (85.3 ft) below top of upper member. Three calcitic specimens of *A.* cf. *A. typicum* Sando in limestone.

● USGS 28604-PC, structurally complex outcrops seemingly near top of lower member. Five calcitic specimens of *A.* cf. *A. typicum* Sando in limestone.

Lithology: Collected sample USGS 29305-PC is dark, partly dolomitized crinoidal grainstone. Collected sample USGS 28604-PC is light, sheared and veined, crinoidal packstone.

Age: The lowest *Ankhelasma* found in this sequence (USGS 28604-PC) is from structurally complex outcrops that seem to belong to the uppermost part of the lower member of the Mission Canyon Limestone. This sample contains *Axinolasma*-like diagnostic corals (*Vesiculophyllum* sp., *Amplexizaphrentis* sp.) and microfossils of Foraminifer Zone 11 or 12 (table A1).

A higher occurrence near the top of the upper member of the Mission Canyon Limestone (USGS 29305-PC) is well located stratigraphically and is well documented by corals of Zone IIIA (*Siphonodendron ocalinum* Sando) both above it (USGS 29306-PC) and below it (USGS 19304-PC). *Ankhelasma* here occurs with microfossils of Foraminifer Zone 11 or 12 (table A1) (USGS 29305-PC), and the occurrence is not far below Foraminifer Zone 12 microfossils (USGS 29309-PC) at the top of the upper member. The total range of *Ankhelasma* at this locality would seem to be from a level low in Composite Zone 13 to about the middle of that composite zone.

12. Little Blackfoot River (Hope Creek) (see Bierwagen, 1964, for description of the locality).

Location: NW¼ SW¼ sec. 29, T. 11 N., R. 6 W., Helena quadrangle (15 min), Lewis and Clark County, Montana.

Stratigraphic position: Mission Canyon Limestone, near top.

Collection: ● USGS 21483-PC, near top of Mission Canyon Limestone. One calcitic specimen of *A.* cf. *A. typicum* Sando in limestone.

Lithology: Collected sample is dark, medium-grained, crinoidal oolitic grainstone.

Age: Approximate stratigraphic position of this occurrence was established by reconnaissance of the outcrop area by Sando. The collected sample contains microfossils of Foraminifer Zone 11 (table A1), which indicates a position in Composite Zone 13.

13. Hannan Gulch (see Mudge and others, 1962, and Mudge, 1972, for description of stratigraphic section).

Location: Sec. 35, T. 22 N., R. 9 W., Sawtooth Ridge quadrangle (7.5 min), Sawtooth Range, Teton County, Montana.

Stratigraphic position: Sun River Member of Castle Reef Dolomite, 48.8 m (160 ft) above base of member.

Collections: ● USGS 18005-PC. One dolomitic specimen (five transverse thin sections) of *A.* cf. *A. typicum* Sando in dolomite.

● USGS 19542-PC. Five dolomitic specimens of *A.* cf. *A. typicum* Sando in dolomite.

Lithology: Collected samples are light, medium- to coarse-grained dolomitized crinoidal grainstone or packstone.

Age: *Ankhelasma* occurs with *Zaphriphyllum* sp., which suggests Coral Zone IIIA for the occurrence. Nichols (1986) reported *Gnathodus texanus* Zone conodonts from the same beds. The zonal level is probably the lower half of Composite Zone 13.

APPENDIX B. CONTROL POINTS FOR CHRONOMETRIC LITHOFACIES PROFILES ACROSS KNOWN VISEAN SHELF MARGIN IN THE UNITED STATES (FIG. 8)

[Interpretations of the stratigraphy by the writers may differ from published references].

Profile *A–A'*

Kane Springs Wash: T. 11 S., R. 64 E., Lincoln County, Nevada. Graphic section by Langenheim (1962, fig. 2, and 1963, fig. 2). Ages determined by physical correlation to Alamo section in T. 6 S., R. 60 E. (Poole and Sandberg, 1977, fig. 2B, loc. 20).

Tungsten Gap: Secs. 11 and 12, T. 14 S., R. 64 E., Clark County, Nevada. Unpublished section measured by Sandberg and Sando. Ages determined by conodonts and corals.

Profile *B–B'*

Dog Valley Peak: Sec. 17, T. 25 S., R. 6 W., Millard County, Utah. Unpublished section measured by Sandberg, Gutschick, and Sando. Part of section described by Sandberg and Gutschick (1984, fig. 10). Ages determined by foraminifers, conodonts, and corals.

Texaco No. 1 Thousand Lake Mountain well: Sec. 11, T. 28 S., R. 4 E., Wayne County, Utah. Well log by Parker and Roberts (1963, fig. 14). Ages determined by physical correlation to Beaverdam Mountains in sec. 35, T. 42 S., R. 18 W., Washington County (Langenheim, 1963, fig. 1).

Profile *C–C'*

City Creek Canyon: Sec. 5, T. 1 N., R. 2 E., Summit County, Utah. Unpublished section measured by Gutschick. Ages determined by corals.

Samak: Sec. 25, T. 2 S., R. 6 E., Summit County, Utah. Section described by Dockal (1980, p. 263–280). Ages determined by corals, foraminifers, and conodonts.

Profile *D–D'*

Old Laketown Canyon: Sec. 32, T. 32 N., R. 6 E., Rich County, Utah. Section described by Sando and others (1959, p. 2762–2763, fig. 5) and reinterpreted by Sando and others (1981, fig. 6, col. 3). Ages determined by corals, conodonts, and foraminifers.

Brazer Canyon: Sec. 20, T. 11 N., R. 8 E., Rich County, Utah. Section described by Sando and others (1959, p. 2751–2754, fig. 5) and reinterpreted by Sando and others (1981, fig. 6, col. 5). Ages determined by corals, conodonts, and foraminifers.

Profile *E–E'*

Wolverine Canyon: Secs. 1, 2, 10, and 11, T. 2 S., R. 38 E., Bingham County, Idaho. Unpublished section measured by Sandberg and Sando. Ages determined by corals and foraminifers.

Fall Creek: Sec. 8, T. 1 N., R. 43 E., Bonneville County, Idaho. Locality studied by Sando and Sandberg. Ages determined by corals.

Sheep Creek: E½ sec. 28, T. 1 N., R. 45 E., Bonneville County, Idaho. Section described by Sando and Dutro (1981). Ages determined by corals and foraminifers.

Profile *F–F'*

Chamberlain Creek: Sec. 20, T. 14 N., R. 29 E., Lemhi County, Idaho. Locality mapped by Skipp and others (1984). Ages determined by conodonts and by physical correlation to Copper Mountain section in secs. 25 and 26, T. 10 N., R. 30 E., Clark County (Huh, 1967, fig. 3, and 1968, section log).

Big Sheep Creek: Sec. 36, T. 14 S., R. 10 W., Beaverhead County, Montana. Unpublished section measured by Sandberg and Sando. Ages determined by corals and foraminifers.

APPENDIX C. *ANAHELASMA* LOCALITIES IN EAST-CENTRAL BRITISH COLUMBIA, CANADA

1. Mountain Creek

Location: Lat 55°23'30" N., long 122°26'00" W.; approximately 2.4 km (1.5 mi) east of Mountain Creek, Pine Pass 1:250,000-scale map sheet (NTS 93O).

Stratigraphic position: Prophet Formation, 129.8–150 m (426–492 ft) below top, upper part of member B.

Collections: ● GSC loc. C4562, 149.3–150 m (490–492 ft) below top of Prophet Formation. Approximately 250 silicified specimens of *A.* n. sp.

● GSC loc. C4563, 135.3–141.4 m (444–464 ft) below top of Prophet Formation. One immature, silicified specimen of *A. typicum*? Sando.

● GSC loc. C4564, 129.8 m (426 ft) below top of Prophet Formation. Two silicified specimens of *A.* sp. indet.

Lithology: Cherty, peloid-bearing, partly dolomitic, spicular, mixed-skeletal packstone and subordinate wackestone and grainstone. Lower part of unit contains dolomitic spiculite.

Collected samples: ● GSC loc. C4562, slightly cherty and argillaceous, pelmatozoan-spicule packstone having corals, brachiopods, and bryozoans.

● GSC loc. C4563, sparsely fossiliferous, finely crystalline dolostone having brachiopods and corals.

● GSC loc. C4564, cherty, peloid-spicule lime packstone having corals, pelmatozoan ossicles, and brachiopods.

Age: All *Ankhelasma* occurrences at this locality are assigned to Composite Zone 14. Associated corals (GSC loc. C4562) are *Aspidiscophyllia*? sp. and *Lophophyllum*? sp., which are zonally nondiagnostic, but microfossils from 19.5 m (63 ft) above *Ankhelasma* belong to Foraminifer Zone 12 (GSC loc. C69462), and those from 11.5 m (37 ft) below *Ankhelasma* are tentatively assigned to Foraminifer Zone 11 or 12 (GSC loc. C69455; table C1).

2. South Mountain Creek

Location: Lat 55°22'00" N., long 122°23'15" W.; approximately 4.2 km (2 mi) east of Mountain Creek, Pine Pass 1:250,000-scale map sheet (NTS 93O).

Stratigraphic position: Prophet Formation, 134.1 m (440 ft) below top, uppermost part of member B.

Collection: ● GSC loc. C110910. Twenty-five silicified specimens of *A.* n. sp.

Lithology: Cherty, spicular limestone similar to the rocks described in the Mountain Creek section (locality 1). No matrix was preserved in the collected sample.

Age: No zonal fossils were found at South Mountain Creek, but in the nearby Mountain Creek section (fig. 12, locality 1), microfossils from approximately the same stratigraphic level belong to Foraminifer Zone 12. Hence, the South Mountain Creek occurrence of *Ankhelasma* is regarded as Composite Zone 14.

3. Sukunka River

Location: Lat 55°01'20" N., long 121°38'25" W.; approximately 6.4 km (4 mi) east of the upper Sukunka River, Dawson Creek 1:250,000-scale map sheet (NTS 93P).

Stratigraphic position: Turner Valley Formation, 7.5 m (24 ft) below top.

Collection: ● GSC loc. C4629. Two silicified specimens of *A.* sp. indet.

Lithology: Packstone and wackestone rich in algae and subordinate echinoderm-bryozoan grainstone in upper part of unit. Main bioclasts are calcareous algae, foraminifers, pelmatozoan ossicles, and ostracodes. Collected sample is peloid-pelmatozoan grainstone having calcareous algae, pellets(?), calcispheres, and foraminifers.

Age: Associated corals are *Zaphrentites* cf. *Z. disruptum* Sutherland, which suggests the upper part of Coral Zone IIIA. Microfossils of Foraminifer Zone 11 (GSC loc. C69613; table C1) occur 11.3 m (37 ft) below *Ankhelasma*. The occurrence of *Ankhelasma* here is regarded as Composite Zone 13.

4. Hook Creek

Location: Lat 54°48'40" N., long 121°39'00" W.; approximately 1.6 km (1 mi) northeast of Hook Lake, Monkman Pass 1:250,000-scale map sheet (NTS 93I).

Stratigraphic position: Turner Valley Formation, 43 m (141 ft) above base.

Collection: ● GSC loc. C7316. Four silicified, immature specimens of *A. typicum*? Sando.

Lithology: Cherty, peloid- and pelletoid-bearing, mixed-skeletal lime grainstone and packstone and subordinate lime wackestone and dolostone. Main bioclasts are calcareous algae, foraminifers, and pelmatozoan ossicles. Collected sample is slightly cherty, pelletoid-foraminifer-pelmatozoan packstone having corals, brachiopods, and peloids.

Age: Associated corals are *Zaphrentites* sp., *Amplexizaphrentis* sp., and *Diastophyllum* cf. *D. inclinatum* Parks, suggestive of Coral Zone IIIA or B. Microfossils from the same stratigraphic level are tentatively assigned to Foraminifer Zone 11 (GSC loc. C90132) (Mamet and others, 1986, p. 43). Microfossils 11.9 m (39 ft) above *Ankhelasma* (GSC loc. C90134) are definitely assigned to Foraminifer Zone 11, and those 25 m (82 ft) below *Ankhelasma* (GSC loc. C90128) belong to the basal part of Foraminifer Zone 10 (Mamet and others, 1986, p. 43). Hence, we regard the occurrence of *Ankhelasma* at this locality as in the lower part of Composite Zone 13.

5. East Fellie Creek

Location: Lat 54°42'50" N., long 120°54'30" W.; approximately 6.4 km (4 mi) northwest of Bone Mountain, Monkman Pass 1:250,000-scale map sheet (NTS 93I).

Stratigraphic position: Turner Valley Formation, 3–4 m (10–46 ft) above base.

Collections: ● GSC loc. C7347, 3 m (10 ft) above base of Turner Valley Formation. Six silicified specimens of *A. typicum* Sando.

● GSC loc. C7350, 14 m (46 ft) above base of Turner Valley Formation. Two silicified, immature specimens of *A.* sp. indet.

● GSC loc. C7351, 13.9 m (45.5 ft) above base of Turner Valley Formation. Two silicified, immature specimens of *A.* sp. indet.

Lithology: Cherty, skeletal dolomite grainstone and fossiliferous, fine- to medium-crystalline dolostone.

Table C1. Foraminifers and algae in samples from Canada Creek in appendix C

(All samples registered in Geological Survey of Canada Locality Catalog. Identifications by Louise Yu, R.J. Mamet)

Taxa	Mountain Creek		Sukunka River	Wapiti River	Jarvis Lake	
	C6882	C6883	C6611	C64248	C7352	C7332
Bisacia sp.	X			X		
Calcitornella sacrul Williamson				X		
C. stachyoides (Pronina)	X	X	X	X		X
C. sp.	X		X	X		
Moravella sp.	X	X		X		
Koninckopora inopinata (Rauser-Chernousova and Reitlinger)			X			
K. sp.				X		
Earlandinella sp.		X		X		X
Baisalina sp.	X			X	X	
Eotuberitina reitlingerae Mamet						
E. reitlingerae (Mamet)		X		X		X
E. sinuosa (Zeller)				X	X	
E. of the group *E. reitlingerae* (Zeller)	X	X	X	X		
E. sp.			X			
Tolypella sp.				X		
Tuberitina collosa Vdovenko				X		
T. sp.				X		
Asteroarchaeolithon hedleyi Mamet			X			
A. sp.	X	X	X	X	X	X
Ivanovella sp.				X		
Komia sp.				X		
Labrinella sp.			X			
Lituotubella sp.				X		
Palaeotextularia sp.				X		
Parachaetetes asbestiformis Komatsu				X		
Parachaetetes sp.	X			X		
Planoendothyra sp.	X					
Pokiella sp.			X			
Priscella sp.	X			X		
Pseudoseptaglomospira sp.				X		
Pseudotextularia sp.				X		
Quasiendothyra sp.				X		
"*Schubertella*" *sensu* Wagner, Hussaini, and Gutschick	X					
Stacheia sp.	X			X		X
Syzrania sp.			X			
Tuberitina sp.			X			

Collected samples • GSC loc. C7347, slightly cherty, dolomitic, pelletoid-bearing, mixed-skeletal wackestone having corals, brachiopods, spicules, calcareous algae, foraminifers, and pelmatozoan ossicles.

• GSC loc. C7350, slightly cherty and dolomitic, pelletoid-pelmatozoan-algal lime packstone having brachiopods.

• GSC loc. C7351, slightly cherty, dolomitic, pelletoid-bearing, mixed-skeletal wackestone having corals, calcareous algae, foraminifers, pelmatozoan ossicles, and pelloids.

Age Associated corals are *Siphonodendron* cf. *S. ovalis*um Sando, *S.* sp., *Vesiculophyllum* sp., *Zaphrentites* sp., and *Syringopora* sp., which indicate Coral Zone IIIA. Associated microfossils (GSC locs. C39452 and C39454) are assigned to Foraminifer Zone 10 (Mamet and others, 1986, p. 45). *Ankhelasma* ranges through the lower one-sixth of Foraminifer Zone 10 here. Microfossils of Foraminifer Zone 9 occur in GSC loc. C39450, 5.8 m (19 ft) below the lowest occurrence of *Ankhelasma* here (Mamet and others, 1986, p. 45). This evidence indicates that *Ankhelasma* is restricted to the lower part of Composite Zone 12 at this locality.

6. Mount Becker

Location Lat 54°31'25" N., long 120°39'45" W., approximately 1.6 km (1 mi) southwest of the peak of Mount Becker, Monkman Pass 1:250,000-scale map sheet (NTS 93I).

Stratigraphic position Mount Head Formation, 47.2–47.9 m (155–157 ft) above base of formation and 0.6–1.2 m (2–4 ft) above base of Baril Member.

Collection • GSC loc. C7377, One silicified, immature specimen of *A.* sp. indet.

Lithology Cherty, pelletoid-bearing foraminifer-pelmatozoan lime grainstone having brachiopod and mollusk fragments. Collected sample is slightly cherty and dolomitic foraminifer-pelmatozoan grainstone having brachiopod and mollusk fragments and pelletoids.

Age Associated corals are *Zaphrentites* sp., which is compatible with Coral Zone IIIB. Microfossils of Foraminifer Zone 12 (GSC loc. C39450) occur 7.6 m (25 ft) below *Ankhelasma* here (Mamet and others, 1986, p. 49). Therefore, the *Ankhelasma* occurrence at this locality is regarded as Composite Zone 13.

7. Wapiti River

Location Lat 54°29'05" N., long 120°43'00" W., approximately 4.2 km (2 mi) east of upper Wapiti River and 8 km (5 mi) northeast of Wapiti Pass, Monkman Pass 1:250,000-scale map sheet (NTS 93I).

Stratigraphic position Mount Head Formation, lower part of Baril Member.

Collections • GSC loc. C64248, lower 3.7 m (5.8 ft) of Baril Member. Ten silicified specimens of *A.* sp. indet.

• GSC loc. C110912, lower part of Baril Member, 85.3 m (280 ft) below top of Mount Head Formation. Four silicified specimens of *A.* sp. indet.

Lithology Pelletoid-bearing, mixed-skeletal grainstone and subordinate fossiliferous dolostone, wackestone, and packstone. Principal bioclasts are corals, calcareous algae, brachiopods, foraminifers, and pelmatozoan ossicles. Some beds cherty. Collected sample (GSC loc. C64248) is pelletoid-bearing, mixed-skeletal grainstone having corals, brachiopods, bryozoans, foraminifers, intraclasts, and pelmatozoan ossicles, and calcisphere-peloid lime wackestone having foraminifers and spicules.

Age Associated corals are *Siphonodendron* sp., *Zaphrentites* sp., *Vesiculophyllum* sp., and *Ekvasophyllum* sp., indicative of Coral Zone IIIA or B. Microfossils in GSC loc. C64248 (table C1) are assigned to Foraminifer Zone 11 or basal 12. Hence, *Ankhelasma* occurs in Composite Zone 13 or the lower part of Composite Zone 14 at this locality.

8. Belcourt Creek

Location Lat 54°21'30" N., long 120°29'40" W., approximately 6.4 km (4 mi) northwest of Mulnok Mountain at head of tributary to Belcourt Creek, Monkman Pass 1:250,000-scale map sheet (NTS 93I).

Stratigraphic position Mount Head Formation, 31.1–55.2 m (102–181 ft) above base (Wileman? and Baril(?) Members).

Collections • GSC loc. C7399, 31.1 m (102 ft) above base of Wileman(?) Member. Two silicified specimens of *A.* sp. indet.

• GSC locs. C7398 and C64249, 15 m (49 ft) above base of Baril(?) Member. Seventeen silicified specimens of *A. typicum* Sando.

• GSC loc. C70699, 38.4 m (126 ft) above base of Wileman(?) Member. Thirty-eight silicified specimens of *A.* sp. indet.

Lithology The Wileman(?) Member is slightly dolomitic, peloid- and pelletoid-bearing, mixed-skeletal wackestone and subordinate shale and marlstone. Principal bioclasts are ostracodes, calcispheres, calcareous algae, molluscs, and pelmatozoan ossicles.

The Baril(?) Member is mainly cherty, skeletal, dolomite grainstone and fossiliferous, medium- to coarse-crystalline dolostone and subordinate cherty, sparsely fossiliferous, fine crystalline dolostone (dolomitized skeletal wackestone?). The main identifiable allochems are pelmatozoan ossicles and silicified corals.

Collected samples • GSC loc. C7399, foraminifer-peloid wackestone having calcispheres and pelmatozoan ossicles.

• GSC loc. C7398, cherty, fossiliferous, fine- to medium-crystalline dolostone having foraminifers, corals, and pelmatozoan ossicles.

• GSC loc. C64249, cherty, fossiliferous, medium-crystalline dolostone having corals and pelmatozoan ossicles.

Age Associated corals are *Zaphriphyllum* sp., *Vesiculophyllum* sp., *Lophophyllum*? sp., and *Syringopora* sp., which suggest Coral Zone IIIA. Microfossils of

Foraminifer Zone 11, GSC loc. C-9025 (1 m/at or the same stratigraphic level as GSC loc. C-7360 and are also present GSC loc. C-9029 (45 ft below GSC loc. C-7358 and C-84249 (Mamet and others, 1986, p. 521). This evidence indicates a Composite Zone 3 occurrence for *Anthracoma* here.

9. Narraway River

Location: Lat 54°10'15" N., long 120°15'20" W., near north of Mount Bannigan, approximately 2.5 km (1.5 mi) southwest of Narraway River, Monkman Pass 1:250,000-scale map sheet (NTS 93I).

Stratigraphic position: 6.1 m (20 ft) above base of Turner Valley Formation to middle? part of Mount Head Formation (faulted section).

Collections:
- GSC loc. C-79613, 6.1 m (20 ft) above base of Turner Valley Formation. Five silicified, immature specimens of *A. typicum*? Sando.
- GSC loc. C-79627, 64.5 m (277 ft) above base of Mount Head Formation (faulted section). One silicified specimen of *A. sp. indet.*
- GSC loc. C-79632, 87 (28.2 m; 414 ft) above base of Mount Head Formation (faulted sections). One silicified specimen of *A. typicum*? Sando.
- GSC loc. C-110911, upper part of Mount Head Formation (faulted section). Four silicified specimens of *A. typicum*? Sando.

Lithology: Alternating succession of dolomitized, skeletal grainstone and sparsely fossiliferous, cherty, fine-crystalline dolomite and subordinate shale and dolomitic marlstone.

Collected samples:
- GSC loc. C-79613, slightly cherty, pellet-bearing, mixed-skeletal grainstone having foraminifers, bryozoans, and pelmatozoan ossicles.
- GSC loc. C-79627, cherty, fossiliferous, medium-crystalline dolostone having pelmatozoan ossicles and corals.
- GSC loc. C-79632, dolomitic, coral-pelmatozoan-spicule packstone.

Age: Associated corals are *Vesiculophyllum* sp., *Zaphrentites* sp., and *Siphonodendron* sp., which suggest Coral Zone IIIA or B. The lowest occurrence of *Anthracoma* (GSC loc. C-79613) is 44 ft below microfossils (GSC loc. C-79617) assigned to Foraminifer Zone 10 by Mamet and others (1986, p. 55, 56) and 186 ft above those (GSC loc. C-79646) of Foraminifer Zone 9 (Mamet and others, 1986, p. 55). Specimens of *Anthracoma* in GSC locs. C-79627, C-79632, and C-110911 occur but cannot be using other fossils but are tentatively assigned to Foraminifer Zone 11 or 12 because of their stratigraphic position several hundred feet above GSC loc. C-79613. *Anthracoma* certainly occurs in Composite Zone 12 and probably ranges into Composite Zone 13 or 14 at this locality.

10. Jarvis Lakes

Location: Lat 54°06'45" N., long 120°13'15" W., approximately 2.4 km (1.5 mi) north of Jarvis Lakes, Monkman Pass 1:250,000-scale map sheet (NTS 93I).

Stratigraphic position: Mount Head Formation, 7.3–43.6 m (24–143 ft) above base.

Collections:
- GSC loc. C-7421, Mount Head Formation, 40.8–43.6 m (134–143 ft) above base. Four silicified specimens of *A. typicum* Sando.
- GSC loc. C-7422, Mount Head Formation, 35.4–36.9 m (116–121 ft) above base. Two silicified, immature specimens of *A. sp. indet.*
- GSC loc. C-7423, Mount Head Formation, 7 m (23 ft) above base. One silicified, immature specimen of *A. sp. indet.*

Lithology: Alternating succession of dolomitized, commonly cherty, skeletal packstone and subordinate, sparsely fossiliferous, fine-crystalline dolomite.

Collected samples:
- GSC loc. C-7421, slightly cherty, pelmatozoan dolomitic grainstone.
- GSC loc. C-7422, brachiopod-bearing, foram-pelmatozoan dolomitic grainstone.
- GSC loc. C-7423, slightly cherty, very dolomitic, pelmatozoan-bearing, mixed-skeletal packstone having calcispheres, bryozoans, foraminifers, and pelmatozoan ossicles.

Age: Associated corals are *Vesiculophyllum* sp., *Amplexizaphrentis* sp., *Lithostrotionella* sp., and *Cyathoclisia*? sp., suggesting Coral Zone IIIA or B. Microfossils (GSC loc. C-9034, table C1 of Foraminifer Zone 11 are associated with the corals in GSC loc. C-7423. Microfossils (GSC loc. C-9032, table C1 of Foraminifer Zone 13 occur more than 150 ft above the highest occurrences of *Anthracoma*, GSC locs. C-7422 and C-7421. The evidence suggests that *Anthracoma* is in Composite Zone 13 and may range into Composite Zone 14 at this locality.

APPENDIX D. CONTROL POINTS FOR CHRONOMETRIC LITHOFACIES PROFILES ACROSS INFERRED VISEAN SHELF MARGIN IN MONTANA (FIG. 13)

(Interpretations of the stratigraphy differ from published references)

Profile *G–G′*

Unnamed, inferred control point, where no record exists, approximately 10 km west of inferred shelf margin.

Rattler Gulch: Secs. 3, 4, and 9, T. 11 N., R. 13 W., Granite County, Montana. Unpublished preliminary study of section by Sando, Perry, and Schneider. Ages determined by corals, conodonts, and foraminifers.

Morris Creek: Secs. 35 and 22, T. 11 N., R. 12 W., Granite County, Montana. Unpublished preliminary study of section by Sando and Perry. Ages determined by corals.

Little Blackfoot River (Hope Creek): Sec. 29, T. 11 N., R. 6 W., Lewis and Clark County, Montana. Unpublished preliminary study of section by Sando and Perry. Locality described by Bierwagen (1964). Ages determined by corals and foraminifers.

Profile *H–H′*

Unnamed, inferred control point, where no record exists, approximately 10 km west of inferred shelf margin.

Lone Butte: Sec. 25, T. 25 N., R. 13 W., Flathead County, Montana. Graphic section by Sloss and Laird (1945, col. 3). Ages determined by brachiopods and corals.

Hannan Gulch: Sec. 18, T. 22 N., R. 9 W., Teton County, Montana. Section described by Mudge and others (1962, fig. 3) and Mudge (1972, p. A97–A98). Ages determined by brachiopods and corals.

Chapter C

Lithostratigraphy and Conodont Biostratigraphy Across the Lower-Middle Ordovician Disconformity (Early to Middle Whiterockian) at Pratt Ferry, Central Alabama

By T.H. SHAW, KEITH E. ROBERSON, and ANITA G. HARRIS

Description and conodont correlation of a stratigraphic unit that formed in paleokarst depressions at the onset of Middle Ordovician transgression

U.S. GEOLOGICAL SURVEY BULLETIN 1895
SHORTER CONTRIBUTIONS TO PALEONTOLOGY AND STRATIGRAPHY

CONTENTS

Abstract C1
Introduction C1
Acknowledgments C2
Location and Procedures C2
Previous Work C3
Lithostratigraphy C3
Newala Limestone C3
Surface of the Lower-Middle Ordovician Disconformity C3
Lenoir Limestone C3
Cottingham Creek Member C3
Mosheim Member C5
Conodont Biostratigraphy C5
Correlation of the Cottingham Creek Member C9
Regional Distribution of the Cottingham Creek Member C10
Conclusions C11
References Cited C12

PLATE

(Plate follows References Cited)

1. Selected conodonts from below and above the Lower-Middle Ordovician disconformity at Pratt Ferry, central Alabama

FIGURES

1. Index map of central Alabama showing distribution of the eastern and western carbonate facies of the Middle and Upper Ordovician C2
2. Chart showing historical development of stratigraphic nomenclature and correlation of some Upper Cambrian to Middle Ordovician rocks in central Alabama C4
3. Photograph and diagrammatic sketch of the type section of the Cottingham Creek Member of the Lenoir Limestone showing location of conodont samples C5
4. Photographs showing karstification features and bedforms at the Lower-Middle Ordovician disconformity, Pratt Ferry, Ala. C6
5. Diagram showing stratigraphic ranges of selected conodont species in the upper part of the Newala Limestone through the lower part of the Mosheim Member of the Lenoir Limestone at Pratt Ferry C8
6. Block diagram showing inferred paleogeographic setting during deposition of the Cottingham Creek Member of the Lenoir Limestone C11

TABLE

1. Stratigraphic distribution of conodont species across the Lower-Middle Ordovician disconformity, Pratt Ferry, Ala. C7

Lithostratigraphy and Conodont Biostratigraphy Across the Lower-Middle Ordovician Disconformity (Early to Middle Whiterockian) at Pratt Ferry, Central Alabama

By T.H. Shaw,[1] Keith E. Roberson,[2] *and* Anita G. Harris[3]

Abstract

Biostratigraphic and lithostratigraphic data from the Newala Limestone (Lower Ordovician) and the disconformably overlying Lenoir Limestone (Middle Ordovician) at Pratt Ferry, Alabama, indicate that (1) the upper Newala Limestone is of late Early Ordovician (late Canadian, lower *Oepikodus communis* Zone) age and the lower Lenoir Limestone is of early Middle Ordovician (late middle to early late Whiterockian; *Phragmodus* "*pre-flexuosus*" Zone to *Cahabagnathus friendsvillensis* Zone) age and (2) a basal, transgressive unit composed primarily of sandy dolostone, the Cottingham Creek Member of the Lenoir Limestone (new name), is the basal unit of the Middle Ordovician sequence locally. Where present, the Cottingham Creek Member aids in the identification of the Lower-Middle Ordovician disconformity in the eastern facies of the Middle Ordovician of Alabama.

The Cottingham Creek Member was deposited in paleokarst depressions developed on the Newala Limestone during the earliest Middle Ordovician. Areal distribution and lithostratigraphic and biostratigraphic data prove conclusively that the Cottingham Creek Member is geographically, lithologically, and chronologically distinct from the Attalla Chert Conglomerate Member of the Chickamauga Limestone of northeastern Alabama, a unit that also lies at the base of the Middle Ordovician transgressive sequence. The Cottingham Creek does appear to be correlative or partly correlative, however, with some of the other units that lie at the base of the Middle Ordovician sequence in the southern Appalachians. It is correlative with the basal part of the Wells Creek Dolomite in south-central Tennessee and the Douglas Lake Member of the Lenoir Limestone in northeastern Tennessee. It may be the same age as the lower part of the Pond Spring Formation in northwestern Georgia but is considerably older than the upper part of the Pond Spring, indicating the possibility of a disconformity within the Pond Spring.

Manuscript approved for publication April 27, 1989

[1] Department of Geology, Queens College, City University of New York, Flushing, NY 11367

[2] Department of Geological Sciences, University of Tennessee, Knoxville, TN 37996

[3] U.S. Geological Survey, 970 National Center, Reston, VA 22092

INTRODUCTION

Recognition of the contact between the Newala Limestone (Lower Ordovician) and the disconformably overlying Lenoir Limestone (Middle Ordovician), which occur together only east of the Helena fault, has long been a problem for geologists mapping in Alabama. This contact, the regionally extensive Lower-Middle Ordovician disconformity, is difficult to locate because of the (1) striking similarity between the Newala Limestone and the Mosheim Member of the Lenoir Limestone, the units that, in most places, bound the disconformity, (2) lack of appreciable relief and other reliable exposure indicators on the surface of disconformity (Roberson, 1988), and (3) generally poor exposure of this stratigraphic interval. The disconformity, however, is well exposed near Pratt Ferry, Bibb County, Ala. (fig. 1), where it is clearly delineated by a large paleokarst collapse feature containing conglomeratic dolostone that differs significantly from either the Newala or Mosheim limestones.

On the basis of new data (Shaw, 1987; Roberson, 1988), this report describes the stratigraphy across the Lower-Middle Ordovician disconformity at Pratt Ferry and establishes the age of the basal dolostone of the Middle Ordovician carbonate sequence, which is named herein the Cottingham Creek Member of the Lenoir Limestone. In addition, the age and origin of the Cottingham Creek Member are compared to those of units occupying a similar stratigraphic position elsewhere in Alabama and in Georgia and Tennessee. The Cottingham Creek Member is inferred to have been deposited in paleokarst depressions on the post-Knox erosion surface during the late middle to early late Whiterockian. This report also establishes the age of the upper Newala Limestone at Pratt Ferry as late Early Ordovician (lower *Oepikodus communis* Zone) and discusses the magnitude of the post-Knox hiatus and the nature of the disconformity surface.



Figure 1. Distribution of the eastern and western carbonate facies in the Middle and Upper Ordovician in central Alabama. The type section of the Cuttingham Creek Member of the Lenoir Limestone is at Pratt Ferry (lower left). Modified from Neathery and Drahovzal, 1986.

Acknowledgments

Portions of this investigation were undertaken as parts of master's theses by T.H. Shaw at Emory University and K.E. Roberson at the University of Alabama, Tuscaloosa. Shaw gratefully acknowledges the aid of his advisors, Pamela Gore and J.E. Repetski. ARCO Oil and Gas Company provided financial support for part of his work; David Hooks, formerly of the University of Alabama, helped him sample the section at Pratt Ferry, and the Illinois State Geological Survey provided facilities and some financial support during preparation of the manuscript. Roberson is indebted to his advisors, D.J. Benson, W.G. Hooks, and C.W. Stock, for sharing with him their knowledge of the Ordovician stratigraphy of Alabama. Some of Roberson's investigations were carried out under the auspices of the Geological Survey of Alabama, and he appreciates that its personnel, particularly G.M. Guthrie, Robert Mink, W.E. Osborne, Dorothy Raymond, W.E. Smith, and B.N. Tew, were readily accessible for discussions. Roberson also wishes to acknowledge financial support provided by Sigma Xi and the Southeastern Section of the Geological Society of America. Special thanks are due J.E. Repetski, U.S. Geological Survey, for sharing his knowledge of Early Ordovician conodonts. We all thank Barbara Stiff, Illinois State Geological Survey, for drafting most of the illustrations.

LOCATION AND PROCEDURES

The exposure near Pratt Ferry is on the east outbank of the Cahaba River, north of the Pratt Ferry Bridge, 0.32 km north of the confluence of Cottingham Creek and the Cahaba River, NE¼NE¼ sec. 33, T. 24 N., R. 10 E., lat 33°01'25" N., long 87°01'18" W., West Blocton East 7.5-minute quadrangle, Bibb County, Ala. (fig. 1). The section is on the north limb of the Bibb County syncline in the Helena thrust sheet.

The section was measured and described, and 74 samples were collected for conodont analysis. Thin sections and acetate peels were made from selected lithologies. The conodont collections are reposited in the U.S. Geological Survey conodont laboratory, Reston, Va.

PREVIOUS WORK

The Lower and Middle Ordovician stratigraphic subdivisions proposed by Butts (1926) have been retained with modification by most workers in Alabama. Butts (1926) recognized that the Upper Cambrian–Lower Ordovician and Middle Ordovician carbonate sequences are separated by an unconformity, and he identified the Mosheim Limestone as the oldest Middle Ordovician unit in the Cahaba Valley east of the Helena fault (fig. 2). The Mosheim is presently considered to be a member of the Lenoir Limestone (Drahovzal and Neathery, 1971; Robertson, 1988; fig. 2).

Drahovzal and Neathery (1971) observed that, although Butts (1926) recognized the facies relations of units within the post-Knox sequence of central Alabama, his nomenclature reflected a time-stratigraphic, succession or "layer-cake" stratigraphy. The general absence of biostratigraphic control and the structural complexity of the Valley and Ridge province resulted in acceptance of Butts' (1926) stratigraphic succession by most subsequent workers. On the basis of lithostratigraphic and paleontologic data, Drahovzal and Neathery (1971) revised the stratigraphic nomenclature to reflect the facies relations of the units recognized by Butts and identified a western, chiefly carbonate facies and an eastern, chiefly clastic facies stratigraphically separated by the Helena fault (figs. 1, 2).

Where the Attalla Chert Conglomerate Member is the lowest unit of the Chickamauga Limestone in the Middle Ordovician western facies, it is the basal transgressive unit of the Middle Ordovician (fig. 2). The Attalla disconformably overlies the Copper Ridge Dolomite, Chepultepec Dolomite, or Longview Limestone and is conformably overlain by the Chickamauga Limestone. The lower Chickamauga in Alabama is late Whiterockian to Blackriveran in age (Hall and others, 1990).

The Attalla Chert Conglomerate Member is absent in the eastern facies of the Middle Ordovician of Alabama, and basal members of the Lenoir Limestone (Middle Ordovician) disconformably overlie the Newala Limestone (upper Lower Ordovician) or Odenville Limestone (uppermost Lower to possibly lowermost Middle Ordovician; fig. 2). Conodonts indicate that the Lenoir is of middle to late Whiterockian age and is time transgressive eastward (Drahovzal and Neathery, 1971; Hall and others, 1990).

LITHOSTRATIGRAPHY

A relatively complete section of lower Paleozoic rocks is exposed in the east outbank of the Cahaba River north of the Pratt Ferry Bridge and includes the Chepultepec, Longview, and Newala Limestones and a nearly complete Middle Ordovician sequence from the base of the Lenoir Limestone through the Athens Shale. The focus of this report is on the disconformable contact between the Newala and Lenoir Limestones.

Newala Limestone

At Pratt Ferry, the Newala Limestone is approximately 85 m thick and consists of medium-bedded, light-gray to very light gray, yellowish-gray-weathering, high-calcium limestone interbedded with thin layers of coarsely crystalline, light brownish-gray-weathering, light-gray dolostone. The limestone is predominantly mudstone and packstone-wackestone. Thin beds of intraclastic debris, grainstone, rudstone, and grainstone are present locally within the lower Newala. With the exception of locally abundant gastropods, megafossils are rare. The Newala conformably overlies the Longview Limestone and is disconformably overlain by the Lenoir Limestone.

Surface of the Lower-Middle Ordovician Disconformity

The disconformable contact between the Newala Limestone and the overlying Cottingham Creek Member of the Lenoir Limestone at Pratt Ferry has a minimum relief of 14 m and forms a paleotopographic low that is at least 48 m across (fig. 3). This depression truncates bedding within the upper Newala. The walls of the depression are very irregular, and the contact between the Newala and rocks filling the depression is sharp (figs. 4A, B). A slightly tilted, downdropped block of Newala Limestone occupies the center of the depression (fig. 3). Relief on the paleokarstic surface away from the depression is low (<10 cm), and the disconformity is represented by a sharp and irregular contact.

The absence of slickensides, fault gouge or breccia, and the continuity of bedding within the depression-fill lithologies and within the Newala on either side of the depression suggest that the depression is the result of karst collapse and not faulting.

Lenoir Limestone

Cottingham Creek Member

The paleotopographic depression in the Newala Limestone is filled with a variety of thin- to medium-bedded, laterally discontinuous, argillaceous, brownish-gray- to light-brownish-gray-weathering, moderate- to light-brown lithologies. These rocks are here named the Cottingham Creek Member of the Lenoir Limestone for Cottingham Creek, Bibb County, Ala. The exposure near

Figure 2. Historical development of stratigraphic nomenclature and correlation of some Upper Cambrian to Middle Ordovician rocks in central Alabama. CCM, shaded unit; Cottingham Creek Member; AC, Attalla Chert Conglomerate Member; OL, Odenville Limestone.

Pratt Ferry, described above, is designated the type section. The base of the member consists of recrystallized conglomeratic dolostone containing pebbles, cobbles, and boulders of lime mudstone, packstone-grainstone, and dolomudstone. The clasts are as much as 30 cm in diameter, and all appear to have been derived from the Newala Limestone (fig. 4C). The basal conglomerate is overlain by dark-brown, bryozoan-rich dolowackestone and dolomudstone; tan, porous dolomudstone; burrowed dolomudstone; and greenish-gray dolosiltite. Crossbedding is well developed



Figure 3. (A) Type section of the Cottingham Creek Member of the Lenoir Limestone and (B) location of conodont samples 5 and 7 to 21 and truncation of bedding within the Newala Limestone and Cottingham Creek Member. Sample numbers are keyed to table 1.

upslip and at the base of the unit (fig. 4D). The uppermost part of the member is a thick-bedded, massive, sandy, argillaceous, brownish-gray dolostone that is laterally continuous across the depression.

Megafossils in the Cottingham Creek Member are predominantly bryozoans but also include brachiopods and large specimens of the gastropod *Maclurites magnus* (LeSueur). The occurrence of these fossils indicates that the depression-fill deposits formed under marine conditions, probably during the earliest phase of the post-Knox transgression.

The maximum thickness of the Cottingham Creek Member is uncertain because the paleotopographic depression that forms its base extends beneath the surface of the Catahba River; it is estimated to be 16 m thick. The Cottingham Creek at Pratt Ferry forms a knoll, possibly because it is composed of dolostone that is relatively more resistant than the adjacent limestones; this topographic characteristic may aid in its recognition elsewhere.

Mosheim Member

The Mosheim Member of the Lenoir Limestone overlies the Cottingham Creek Member. Its lower part is predominantly medium- to thick-bedded, massive, medium-light- to light-gray, fenestral lime mudstone containing a sparse fauna of ostracodes and gastropods. These beds grade upward into thin- to medium-bedded, slightly more argillaceous, fenestral mudstone that locally contains abundant gastropods. The contact between the Mosheim and Cottingham Creek Members is sharp and irregular and may represent an erosional surface; if so, the magnitude of the hiatus is less than a single conodont zone.

CONODONT BIOSTRATIGRAPHY

The rather complete exposure of Ordovician rocks in the vicinity of Pratt Ferry, Ala., has become a key section for biostratigraphic studies in the southernmost Appalachians, particularly for Middle Ordovician index fossil groups (brachiopods: Cooper, 1956; conodonts: Sweet and Bergstrom, 1962, Bergstrom, 1971, and Schmidt, 1982; graptolites: Ruedemann, 1947, Decker, 1952, and Finney, 1977, 1984). In this report, we add to the paleontologic documentation of this section the distribution of conodonts in several meters of beds both below and above the



Figure 4. Karstification features and bedforms at the Lower-Middle Ordovician disconformity, Pratt Ferry, Ala. *A*, Sharp, vertical contact (circled area) between the Newala Limestone (lighter gray) and the Cottingham Creek Member of the Lenoir Limestone. *B*, Close-up of contact circled in *A*. *C*, Boudget of Newala Limestone in the lower Cottingham Creek Member. *D*, Crossbedded dolostone at base of Cottingham Creek Member.

discontinuity separating the Newala and Lenoir Limestones, an interval not covered by previous conodont studies.

Of the 34 samples collected for conodont analysis, 31 yielded conodonts. Sample size ranged from 3.9 to 12.1 kg (table 1). Large samples are required to recover even a few conodont elements from either the Newala Limestone or the lower members of the Lenoir Limestone. The ranges of selected conodont species in the section are shown in figure 5; conodont species distribution is summarized in table 1. Conodonts from this section have a color alteration index of 1.5; this index indicates the host rock reached burial temperatures of at least 80 to 90 °C.

The conodont fauna from the Newala is dominated by elements of *Oneotodus beaverfieldensis* (Branson and Mehl) (pl. 1, figs. 18, 19) and *Eucharodus parallelus* (Branson and Mehl) and subordinate numbers of elements of *Cordylodus lenoirus* Repetski (pl. 1, fig. 21), *Cordylodus caseyi* (Branson and Mehl) (pl. 1, fig. 22), *Drepanoistodus suberectus* (Branson and Mehl) subsp. A *sensu* Repetski, 1982a, *Glyptoconus quadriplicatus* (Branson and Mehl), *Oistodus venustus* Ethington and Brand (pl. 1, fig. 20), *Scolopodus cornuformis* Branson and Mehl, *Scolopodus emarginatus* Barnes and Tuke (pl. 1, fig. 23), *Teridontus nakamurai* Branson and Mehl, and *Utahconus utahensis* Branson and Mehl.

Although *Oepikodus communis* has not been found in the Newala Limestone, probably due to unfavorable environmental conditions, *Diaphorodus delicatus*, a species whose first appearance approximates the base of the *O. communis* Zone in rocks that formed in warm, shallow, possibly hypersaline marine environments (Repetski, 1985), occurs throughout the upper 25 m of the formation in the Pratt Ferry section. *D. delicatus* is known to range into the early part of Whiterockian. Its occurrence in the Newala, together with representatives of *Cordylodus intermedius* and *Glyptoconus quadriplicatus* (table 1), species that are not known to range into the Whiterockian, indicates that the

Table 1. Stratigraphic distribution of conodont species across the Lower-Middle Ordovician disconformity, Pratt Ferry, Ala.



Figure 9. Ranges of selected conodont species in the upper part of the Newala Limestone and lower part of the Lenoir Limestone at the type section of the Cottingham Creek Member of the Lenoir Limestone. Extension of ranges below and above the stratigraphic levels of the Newala and the Lenoir Limestones sampled for this study are for the Newala, from Shaw (1987), and for the Lenoir, from Sweet and Bergström (1962), Bergström (1971), and Schmidt (1982). (A) *Drepanodus concavus*, (B) *Eoplacognathus pseudoplicatus*, (C) *Cryptoconus quadriplicatus*, (D) *Coelocerodontus costatus*, (E) *Unicodina abnormalis*, (F) *Periodon* complex, (G) *Costodus lacunatus*, (H) *Diaphorodus delicatus*, (I) "*Scolopodus*" *emarginatus*, (J) *Amorphognathus* sp., (K) *Belodina monitorensis*, (L) *Plectodina cochleariformis*, (M) "*Scandodus*" *simplex*, (N) *Leptochirognathus* sp., (O) *Protopanderodus* aff. *P. balticus*, (P) *Cahabagnathus friendsvillensis*, and (Q) *Phragmodus flexuosus*. Conodont distribution and abundance are given in table 1.

upper Newala was deposited during *O. communis* Zone time (= late Early Ordovician). Moreover, the presence of *C. lonodon* throughout much of this interval further restricts these beds to the lower part of the *O. communis* Zone.

The conodont fauna of the Cottingham Creek Member of the Lenoir Limestone consists chiefly of, in order of decreasing abundance, elements of *Panderodus* spp. (pl. 1, fig. 17), *Amorphognathus ethingtoni* and Schumacher, *Plectodina cochleariformis* (Andrews) (pl. 1, fig. 15), *Belodina monitorensis* Ethington and Schumacher, *Drepanoistodus* spp., *Erismodulus* aff. *E. balticus* Dzik, and "*Scandodus*" *simplex* Mound. Reworked elements of *Drepanodus concavus*, *Eoplacognathus pseudoplicatus*, *Cryptoconus quadriplicatus*, *Unicodina abnormalis*, and "*Scandodus*" *emarginatus* are locally present within the Cottingham Creek Member.

With the exception of "*Scandodus*" *simplex*, the Middle Ordovician species recovered from the Cottingham Creek Member also occur in the overlying Mosheim Member. In addition, elements of *Tripodus* sp., *Oistodus* cf. *O. lanceolatus* Pander, *Drepanoistodus* cf. *D. suberectus* (Lindström), *Cahabagnathus friendsvillensis* (Bergström) (pl. 1, fig. 1), and *Phragmodus flexuosus* Moskalenko (pl. 1, figs. 2–4), first occur in the Mosheim Member.

Specimens of "*Scandodus*" *simplex* are the oldest, possibly indigenous conodonts in the basal beds of the Cottingham Creek Member. If they are indigenous and not reworked, then their occurrence in this member, together with elements of *Plectodina cochleariformis* and *Erismodulus* aff. *E. balticus*, restricts the age of these beds to the *Phragmodus* "*pre-flexuosus*" Zone of Sweet (1984). The upper limit of the range of "*S.*" *simplex* and the lower limit of the ranges of *P. gracilimarginatus* and *E. balticus* appear to be within this chronozone. If, on the other hand, the specimens of "*S.*" *simplex* are reworked (these specimens are limited to beds that contain other reworked conodont elements), then the age of the Cottingham Creek Member is less constrained because all other conodonts within this unit are relatively long ranging within the middle to late Whiterockian. Elements of *Cahabagnathus friendsvillensis* first appear several meters above the base of the Mosheim Member in this exposure (fig. 5), indicating a level within the *C. friendsvillensis* Zone of the North American conodont zonation (Sweet, 1984) or the *P. serra* Zone of the Baltoscandia conodont zonation (Bergström (1971) and oral commun., 1988); recovered elements of *Pygodus serra* Hadding and *C. friendsvillensis* from near the top of the Lenoir to at least 65 m below its top in the vicinity of Pratt Ferry. Bergström (1971) has also shown that the boundary between the *P. serra*–*P. anserinus* Zones lies near the top of the Pratt Ferry Formation (Cooper, 1956), about 10 m above the top of the Lenoir Limestone in this vicinity. The known range of *C. friendsvillensis* is from near the base to near the top of the *P. serra* Zone (Bergström, 1983). Its presence throughout the nearly 75 m (+) Lenoir Limestone that overlies the Cottingham Creek and lower Mosheim beds in the Pratt Ferry section suggests that the Cottingham Creek Member is no younger than the early part of the *C. friendsvillensis* Zone and could well be as old as, but no older than (based on the presence of *Plectodina cochleariformis* and *Erismodulus* aff. *E. balticus*), the *Phragmodus* "*pre-flexuosus*" Zone. In terms of the Whiterockian Series, the Cottingham Creek Member is of late middle to early late Whiterockian age.

CORRELATION OF THE COTTINGHAM CREEK MEMBER

The Cottingham Creek Member of the Lenoir Limestone is differentiated from the Attalla Chert Conglomerate Member of the Chickamauga Limestone by the absence of chert, the sparsity of cobble- and boulder-sized clasts, and the absence of sandstone and tan green shale in the Cottingham Creek Member. Lithologic differences between the Cottingham Creek and Attalla Chert Conglomerate Members are the result of differences in provenance, that is, differences in the rock units that underlie them and the nature of the paleotopographic depressions in which each member was deposited.

On the basis of conodonts in Chickamauga Limestone beds above the Attalla Chert Conglomerate Member at the Red Mountain road cut in Birmingham, Ala., Hall and others (1986) reported the age of the Attalla as early Blackriveran. Their lowest, biostratigraphically useful, Middle Ordovician conodont collection is from about 12 m above the top of the Attalla, and it and overlying collections only indicate that the Attalla is no younger than latest Whiterockian to early Blackriveran. About 20 km east-northeast of Birmingham, we collected one conodont sample from a Middle Ordovician dolomitic terrestrial mudstone, 2 m below the Attalla and another sample from within a noncherty phase of the Attalla (USGS collections 10315-CO and 10316-CO) from outcrops on the northern side of a stream that flows between Butler and Foster Mountains, 400 m east of the Clay-Palmerdale Road and 2 km northwest of Clay, Jefferson County, SE¼NW¼ sec. 23, T. 13 S., R. 1 W., lat 33°42.8′ N., long 86°57.1′ W., Argo 7.5-minute quadrangle, Ala.). The presence of representatives of *Phragmodus undatus* Stauffer and *Plectodina* n. sp. of Votaw (1971) in the lower sample and of these two species and *Aspidagnathus delicatus* Bergström and others in the Attalla Chert Conglomerate Member sample indicates that the Attalla is of highest *C. tricostatus* Zone to lowest *E. gundersoni–dactylus* Zone age. Thus, even though the Cottingham Creek and the Attalla Chert Conglomerate Members formed in similar paleogeographic settings and are the basal units of the Middle Ordovician carbonate sequence that disconformably overlies the Knox Group, they are neither coeval nor contiguous (fig. 2). The Attalla occurs only west of the Helena fault, whereas the Cottingham Creek is found only east of it.

The age relation of the Cottingham Creek Member to the Pond Spring Formation of northwestern Georgia is more complex. The basal few meters of the Pond Spring are lithologically similar to the Cottingham Creek Member and occupy a similar stratigraphic position. In the vicinity of Chickamauga, Ga., its type area, the Pond Spring ranges from about 75 to 95 m in thickness (Miller and Smith, 1960). According to Hall and others (1986), the lowest occurrence of *Plectodina aculeata* (Stauffer) in this area is about 15 m below the top of the formation and about 14 m above the only sample containing *Phragmodus flexuosus* and *P. inflexus*. These data indicate that the lowest sample containing *P. undatus* is within the *P. aculeata* Zone and that at least the upper Pond Spring is of Blackriveran age and, like the Attalla Chert Conglomerate Member, is younger than any part of the Lenoir Limestone at Pratt Ferry. According to Schmidt (1982), two collections from the lower 20 m of the Pond Spring contain conodonts much older than those from the upper part of the formation. These collections contain abundant representatives of *Phragmodus flexuosus*, lesser numbers of its ancestor *P.* "*pre-flexuosus*," and forms transitional between them. These data seem to indicate that at least the lower 20 m of the Pond Spring are of lowest *C. tricostatus* Zone age. This age is somewhat surprising, because the highest occurrence of this species association is only about 30 m below Pond Spring beds that contain *P. aculeata* and that are of definitive Blackriveran age. Moreover, the regional depositional framework and lithology of the Pond Spring suggest rapid deposition. A disconformity may occur within the middle Pond Spring, or alternatively, the specimens assigned by Schmidt (1982) to forms ancestral and transitional to *P. flexuosus* may be ecophenotypes that appear late in the range of *P. flexuosus*. Schmidt (1982) expressed similar concern but offered no suggestions about the apparent long time interval represented by the lower and middle Pond Spring Formation. This problem cannot be resolved without additional data. Nonetheless, at least the lower 20 m of the Pond Spring may be equivalent to the Cottingham Creek Member.

The Cottingham Creek Member is at least partly correlative with the base of the Wells Creek Dolomite (as used by Bentall and Collins, 1945), the unit that disconformably overlies the Knox Group in the subsurface of south-central Tennessee. Conodont collections from the basal few meters of the Wells Creek in four cores from Bedford and Maury Counties (Repetski, 1982b, and unpub. data) contain representatives of *Multioistodus subdentatus* Cullison and *Phragmodus flexuosus*, indicating a level low in the *C. friendsvillensis* Zone. Except for a few redeposited Early Ordovician specimens, the Middle Ordovician conodonts in these collections appear to be indigenous. The

base of the Wells Creek Dolomite in at least Bedford and Maury Counties, Tenn., is the same or nearly the same age as the Cottingham Creek Member and appears to be the same age as the lower part of the Pond Spring Formation of northwestern Georgia.

The Cottingham Creek Member is probably correlative with the Douglas Lake Member of the Lenoir Limestone in Tennessee (Laurence, 1944; Bridge, 1955; Harris, 1982). Like the Cottingham Creek, the Douglas Lake Member formed within paleokarst depressions on the post-Knox erosion surface. Rocks of the Cottingham Creek are similar to rocks assigned to unit B by Laurence (1944), which include cross-bedded conglomeratic dolostone, and recrystallized argillaceous dolostone. The Douglas Lake Member disconformably overlies the Mascot Dolomite, a cherty dolomite that is coeval in part with the Newala Limestone. The Douglas Lake consists predominantly of noncarbonate rocks that contain chert locally and thus are unlike rocks of the Cottingham Creek Member. Differences in rock type reflect the composition of the underlying Knox Group and the paleotopography and paleoregolith developed on the surface of disconformity.

Laurence (1944) reported the age of the Douglas Lake Member as post-Canadian and pre-Chazyan or Chazyan or middle to late Whiterockian of current usage (Ross and others, 1982). Harris (1982) and Brindley and others (in press) reported elements of *Phragmodus flexuosus* from about 8 m above the top of the Douglas Lake Member in the main body of the Lenoir Limestone. In a collection taken 20 m higher in the section from the overlying Whitesburg Formation, they reported the presence of *Eoplacognathus suberectus* (Fahraeus), *Coelocerognathus trombiformis*, and *Pygodus serra*. These data indicate that the Douglas Lake Member is no younger than the lowest part of the *E. suberectus-trombiformis* Zone and is of that age or of slightly older late *P. "pre-flexuosus"* Zone age and falls within the age range determined for the Cottingham Creek Member.

REGIONAL DISTRIBUTION OF THE COTTINGHAM CREEK MEMBER

Rocks resembling and occurring at the same stratigraphic level as the Cottingham Creek Member have been identified near Fremont Gap, St. Clair County, Ala. (SE ¼ sec. 9, T. 14 S., R. 5 E.; Roberson, 1988). At both Pratt Ferry and Fremont Gap, these rocks occur in paleokarst depressions developed on the Newala Limestone. At Fremont Gap, the dolostones are very similar to those at the type section of the Cottingham Creek; however, the basal conglomerates are less conspicuous, and the geometry of the deposit is difficult to determine because of extensive covered intervals.

The Cottingham Creek is apparently restricted to areas where the Newala Limestone is the highest unit preserved beneath the Lower-Middle Ordovician disconformity. This may be related to the character of the Newala, local structure, magnitude of the hiatus, and (or) proximity to the source area for clasts within Cottingham Creek conglomerate.

Paleotopographic depressions resulting from karst collapse or dissolution have not been observed where the Odenville Limestone is the highest unit preserved beneath the post-Knox Group disconformity. Dolomitic, siliceous, and argillaceous impurities within the Odenville, typically 10 to 12 percent, may have inhibited the development of large subsurface solution cavities by retarding migration of ground water. In contrast, the Newala Limestone rarely contains more than two percent noncarbonate impurities and appears to be very susceptible to aqueous dissolution.

Nodular chert-type, coarsely crystalline dolostone within the Newala Limestone may increase its dissolution potential relative to that of the Odenville or Longview Limestones. Rocks of this type are highly susceptible to sinkhole formation (Brook, 1982). Blowzone caverns within the lower Newala are commonly encountered during drilling in the type area of the formation (B. Kitchens, Vulcan Materials Corporation, and T. Flannery, Blue Circle Industries, oral commun., 1988). A collapsed cave occurs within the lower Newala near the base of the Vulcan Materials Corporation quarry north of Calera, Ala.; the age of this feature is unknown. Holocene sinkholes in the type area of the Newala are most common in the adjacent Longview Limestone because this unit contains tectonically fractured chert that provides pathways for fluid migration (LaMoreaux, 1984).

Local structure may also significantly influence the development of relief on the Knox paleokarst surface. The Attalla Chert Conglomerate Member is restricted to an area between the Anniston and Harpersville lineament complexes (fig. 3; Drahovzal, 1974). Moreover, the thickest and coarsest development of the Attalla is associated with the lineament complexes (Drahovzal and others, 1974); within the Anniston lineament complex, high-angle faults vertically offset the top of the Knox Group by as much as 45 m. These faults may be the result of differential movement of crustal blocks contemporaneous with or prior to deposition of the Attalla Chert Conglomerate Member (Drahovzal and others, 1974). These data imply that development of the Attalla is related to regional structural features.

In contrast, the known Cottingham Creek deposits do not appear to be associated with regional structures. At the type section of the Cottingham Creek, there is no evidence for tectonic deformation of the Newala prior to deposition of the Cottingham Creek, and stratigraphy appears to be the controlling factor for paleokarst development.

Duration of exposure may also significantly influence the configuration of the paleokarst surface developed on the Knox Group. Rigenbach and Harris (1976) and Roberson (1988) reported that the magnitude of the post-Knox–Middle Ordovician disconformity generally decreases eastward in the Appalachian basin. In addition, Neathery and Drahovzal (1973, 1988) reported a dark-maroon shale and a red to pink cherty clay stone near the base of the Attalla Chert Conglomerate Member and interpreted these rocks to represent a paleosol developed on the Knox surface. The apparent decrease in magnitude of the post-Knox–Middle Ordovician disconformity eastward and the development of a paleosol at the base of the Attalla imply that rocks below the disconformity in the western facies belt may have been exposed long enough to develop a valley and hill topography having a primarily surficial drainage, whereas rocks of the eastern facies belt may have been submerged before a surficial drainage pattern could be established. Furthermore, tectonic deformation of the chert-bearing rocks underlying the Attalla Chert Conglomerate associated with the Anniston and Harpersville lineaments may have accelerated degradation of the Knox paleosurface and dissolution in the subsurface. The Attalla Chert Conglomerate Member has a greater areal distribution than the Cottingham Creek Member. This supports the hypothesis that the Attalla Chert Conglomerate Member may represent valley-fill deposits rather than locally restricted karst-depression fills representative of the Cottingham Creek deposits.



Figure 6. Interpreted paleogeographic setting during deposition of the Cottingham Creek Member of the Lenoir Limestone in what is now part of central Alabama. NWB, normal wave base.

Neathery and Drahovzal (1973, 1986) reported that the Attalla Chert Conglomerate Member near Hensley Mountain, Ala., exhibits an upward size gradation and an increase in clast size approximately 12 m above the base of the unit. They attributed this increase to renewed influx of clastic material into the basin. Furthermore, some clasts are rounded and suggest mechanical transportation. The absence of conglomerate above the basal beds of the Cottingham Creek Member and the general angularity of clasts in these beds indicate their local derivation.

Figure 6 shows our interpretation of the paleogeographic setting when Cottingham Creek deposits accumulated in karst depressions within the Newala Limestone during the late middle to early late Whiterockian in the area that is now central Alabama. About 30 to 40 km northeast of Pratt Ferry and east of the Helena fault, the Odenville Limestone conformably overlies the Newala Limestone. Conodonts from the Odenville (J.E. Repetski, U.S. Geological Survey, unpub. data) indicate a very latest Early Ordovician age (highest *O. communis* Zone) to possibly very earliest Whiterockian age. This suggests that the depression at Pratt Ferry formed during early or middle Whiterockian time.

CONCLUSIONS

At Pratt Ferry, Ala., the base of the Lenoir Limestone consists of argillaceous dolostone and conglomerate

Because these rocks are unlike other rocks in the Lenoir, they are herein designated the Cottonplant Creek Member of the Lenoir Limestone. Conodonts indicate that this new member is of late middle to early late Ordovician (late and post-Whiterockian) age, and therefore the Nemagin Limestone is of late Early Ordovician age. The contact of the Cottonplant Creek Member and the overlying Mossbenn Member of the Lenoir Limestone is sharp and irregular and represents a disconformity surface of relatively short duration. The Cottonplant Creek deposits occur in paleokarst depressions developed on the Nemagin during the early and (or) middle Whiterockian. Areal distribution and lithostratigraphic and biostratigraphic data prove conclusively that the Cottonplant Creek Member is geographically, lithologically, and chronologically distinct from the Wells Creek Conglomerate Member. The Cottonplant Creek is correlative with the Douglas Lake Member of the Lenoir Limestone of northeastern Tennessee, at least the lowest beds of the Wells Creek Dolomite of south-central Tennessee, and only the lowest part, apparently, of the Powell Spring Formation of northwestern Georgia.

REFERENCES CITED

Bassler, Ray, and Cullison, J.S., 1943, Silurian invertebrates and stratigraphy of the pre-Tuscumbia Outlier area and Giant Cumberland cuesta in central Tennessee: Tennessee Division of Geology, Oil and Gas Investigations Preliminary Chart 4.

Bergström, S.M., 1971, Conodont biostratigraphy of the Middle and Upper Ordovician of Europe and eastern North America, *in* Sweet, W.C., and Bergström, S.M., eds., Symposium on conodont biostratigraphy: Geological Society of America Memoir 127, p. 83–161.

———1983, Biogeography, evolutionary innovations, and mass extinction in Ordovician plankton, *in* Nitecki, M.H., ed., Biotic crises in ecological and evolutionary time: Academic Press, Inc., p. 61–78.

Bergström, S.M., Ethington, R.L., and Ross, R.J., Jr., 1965, Conodonts of the Vinini Formation and their significance for the classification, correlation, and evolution of Ordovician conodont faunas: Journal of Paleontology, v. 39, no. 2, p. 167–201.

Bergström, S.M., Huff, W.D., and Kolata, D.R., 1992, Graptolite biostratigraphy of the Lower Member of the Lexington Limestone (Middle Ordovician) in the subsurface of central Kentucky: Journal of Paleontology, v. 66, no. 5, p. 729–778.

Bergström, S.M., Kolata, D.R., and Huff, W.D., 1983, The Deicke bentonite (Upper Ordovician) in the Appalachian basin: Correlation, sedimentology, and biostratigraphy: Geological Society of America Bulletin, v. 94, p. 1248–1266.

Bergström, S.M., Kolata, D.R., Huff, W.D., and Kleffner, M.A., 1992, Graptolite biostratigraphy of the Lexington Limestone (Middle Ordovician) in the subsurface of central Kentucky: Journal of Paleontology, v. 66, no. 5, p. 779–798.

Bergström, S.M., Kolata, D.R., Kleffner, M.A., and Huff, W.D., 1992, Graptolite biostratigraphy of the Middle Ordovician Carbonate Group and related strata of the Appalachian basin, *in* Bergström, S.M., and Dorobek, S.L., eds., Carbonate facies in the 25th annual field conference of the Appalachian Geological Society: Tuscaloosa, Alabama, Appalachian Geological Society, p. 83–111.

Bergström, S.M., Kolata, D.R., and Kleffner, M.A., 1995, Biostratigraphy and sedimentary significance of conodonts from the Middle Ordovician Lebanon Limestone and Leipers Limestone, Nashville Dome, Tennessee: Journal of Paleontology, v. 69, no. 6, p. 1115–1135.

Bergström, S.M., Kolata, D.R., and Kleffner, M.A., 1995, Correlation of the Middle Ordovician Wardell Formation, Missouri, and the Dutchtown Limestone, Illinois, with the Middle Ordovician Decorah Subgroup, Upper Mississippi Valley: Journal of Paleontology, v. 69, no. 6, p. 1136–1151

PLATE 1

SEM photomicrographs of specimens coated with carbon and gold; all specimens rep
in the U.S. National Museum of Natural History (USNM), Washington, D.C.

Contact photographs of the plates in this report are available, at cost, from the U.S. Geolog
Photographic Library, Federal Center, Denver, CO 80225

PLATE 1

Selected conodonts from below and above the Lower-Middle Ordovician disconformity at Pratt Ferry, central Alabama

Figures 1–13. Conodonts from the Mosheim Member of the Lenoir Limestone

1. *Cahabagnathus friedmanellensis* (Bergström), upper view of pastinplanate element. ×75. USGS colln. 10088-CO, USNM 430386.

2–4. *Phragmodus flexuosus* Moskalenko, lateral views of M, P (outer), and Sc elements. ×75. USGS colln. 10091-CO, USNM 430387–89.

5–7. *Ansella mesakmensis* (Ethington and Schumacher), lateral views of P, M, and Sa elements. ×50. USGS colln. 10086-CO, USNM 430390–92.

8, 9. *Belodina monitorensis* Ethington and Schumacher, inner lateral views of cordodontiform and rostrate elements. ×75. USGS colln. 10091-CO, USNM 430393–94.

10–12. *Erraticodon* aff. *E. balticus* Dzik. ×50.

10. Anterior view of Pb element. USGS colln. 10091-CO, USNM 430395.

11, 12. Inner lateral views of M and Sc (anterolateral process broken) elements. USGS colln. 10086-CO, USNM 430396–97.

13. *Leptochirognathus quadritinis* Bramon and Mehl, inner lateral view of prioniodontiform element. ×75. USGS colln. 10086-CO, USNM 430398.

14–16. *Plectodina aculeata* (Andrews). ×75.

14, 16. Inner lateral views of Pa and Sc elements, Mosheim Member of Lenoir Limestone, USGS colln. 10086-CO, USNM 430399, 430816.

15. Posterior view of Sb element, Cottingham Creek Member of Lenoir Limestone, USGS colln. 10046-CO, USNM 430817.

17. *Panderodus* sp., outer lateral view. ×75. Cottingham Creek Member of Lenoir Limestone, USGS colln. 10085-CO, USNM 430400.

18–23. Conodonts from the upper part of the Nolichucky Limestone.

18, 19. *Diaphorodus delicatus* (Bramon and Mehl), inner lateral views of M and Sb elements. ×75. USGS colln. 10039-CO, USNM 430401–02.

20. *Oneotodus costatus* Ethington and Brand, inner lateral view (anterobasal margin partly broken). ×75. USGS colln. 10080-CO, USNM 430403.

21. *Cristodus loxodus* Repetski, inner lateral view of rostrate element. ×75. USGS colln. 10082-CO, USNM 430404.

22. *Drepanodus concavus* (Bramon and Mehl), inner lateral view of arcuate form element. ×75. USGS colln. 10033-CO, USNM 430405.

23. "*Scolopodus*" *emarginatus* Barnes and Tuke, inner posterolateral view of Sb? element. ×75. USGS colln. 10080-CO, USNM 430406.

CAHABAGNATHUS, PHRAGMODUS, ANSELLA, BELODINA, ERRATICODON, LEPTOCHIROGNATHUS, PLECTODINA, PANDERODUS, DIAPHORODUS, ONEOTODUS, CRISTODUS, DREPANODUS, AND "SCOLOPODUS"

SELECTED SERIES OF U.S. GEOLOGICAL SURVEY PUBLICATIONS

Periodicals

Earthquakes & Volcanoes (issued bimonthly).

Preliminary Determination of Epicenters (issued monthly).

Technical Books and Reports

Professional Papers are mostly comprehensive scientific reports of wide and lasting interest and importance to professional scientists and engineers. Included are reports on the results of resource studies and of topographic, hydrologic, and geologic investigations. They also include collections of related papers addressing different aspects of a single scientific topic.

Bulletins contain significant data and interpretations that are of lasting scientific interest but are generally more limited in scope or geographic coverage than Professional Papers. They include the results of resource studies and of topographic, hydrologic, and geologic investigations. They also include collections of related papers addressing different aspects of a single scientific topic.

Water-Supply Papers are comprehensive reports that present significant interpretations and detailed scientific data on the quantity, quality, and distribution of water and in the hydrologic environment.

Circulars present administrative information or miscellaneous scientific information of wide popular interest in formats designed for distribution at no cost or at low cost to the public.

Water-Resources Investigations Reports are primarily of the open-file variety and are available for consultation at depositories. Some are also available for purchase at inexpensive prices from the U.S. Geological Survey, Books and Open-File Reports Section, Federal Center, Box 25425, Denver, CO 80225. Check the publication catalogs or with the Publications Specialist nearest you.

Open-File Reports include unpublished manuscript reports, maps, and other material that are made available for public consultation at depositories. They are a nonpermanent form of publication that may be cited in other publications as sources of information.

Maps

Geologic Quadrangle Maps are multicolored geologic maps on topographic quadrangle bases at various scales (commonly 1:24,000 or 1:62,500) showing bedrock or surficial geology, stratigraphic and structural relations in addition to lithology. Most quadrangles are mapped at 1:24,000 or 1:62,500 scale, but some are mapped at smaller or larger scales.

Geophysical Investigations Maps are topographic or planimetric maps that portray geophysical data (such as gravity, magnetic, seismic, or radioactivity) and sometimes include geologic interpretations. Most maps are published at the same scale as the topographic base map.

Miscellaneous Investigations Series Maps are published at various scales. Some consist of a single map; others consist of several maps accompanied by explanatory text. Some are folded maps intended for field use; others are wall maps. Many of these maps portray the geology of the Moon.

Coal Investigations Maps are geologic maps on topographic or planimetric bases at various scales showing bedrock or surficial geology, stratigraphy, and structural relations in areas containing coal resources.

Oil and Gas Investigations Charts show stratigraphic information for oil or gas fields and other areas having petroleum or natural-gas potential.

Miscellaneous Field Studies Maps are multicolored or black-and-white maps on topographic or planimetric bases presenting a wide range of geologic, hydrologic, and mineralogic information. Some of these maps are published at the same scale as the base map, but many are published at different scales.

Hydrologic Investigations Atlases are multicolored or black-and-white maps on topographic or planimetric bases presenting a wide range of geohydrologic data of both regional and specialized interest. The scale of these maps is 1:24,000 and smaller.

Catalogs

Periodically, as well as at other times, printing a comprehensive listing of U.S. Geological Survey publications that are available under the conditions indicated below from the U.S. Geological Survey, Books and Open-File Reports Section, Federal Center, Box 25425, Denver, CO 80225. See price and availability list.

"Publications of the Geological Survey, 1879–1961" may be purchased by mail and over the counter in paperback book form and as a set of microfiche.

"Publications of the Geological Survey, 1962–1970" may be purchased by mail and over the counter in paperback book form and as a set of microfiche.

"Publications of the U.S. Geological Survey, 1971–1981" may be purchased by mail and over the counter in paperback book form (two volumes, publication years listed separately) and as a set of microfiche.

Supplements for 1982, 1983, 1984, 1985, 1986, and for subsequent years until the next permanent catalog has been published by mail and over the counter in paperback book form.

State Catalogs, "List of U.S. Geological Survey Geologic and Water-Supply Reports and Maps For (State)," may be purchased by mail and over the counter in paperback booklet form only.

"Price and Availability List of U.S. Geological Survey Publications," issued annually, is available free of charge in paperback booklet form only.

Selected copies of a monthly catalog "New Publications of the U.S. Geological Survey" are available free of charge by mail or may be obtained over the counter in paperback booklet form only. Those wishing a free subscription to the monthly catalog "New Publications of the U.S. Geological Survey" should write to the U.S. Geological Survey, 582 National Center, Reston, VA 22092.

Note. — Prices of Government publications listed in older catalogs, announcements, and publication lists may be incorrect. Therefore, the prices charged may differ from the prices in catalogs, announcements, and publications.